

**Inventor search history**

=> d his L42

(FILE 'HCAPLUS' ENTERED AT 16:41:51 ON 19 OCT 2009)

L42 16 S L41 AND (GUAN? OR BIOFILM? OR BIO(W)FILM? OR "BIOFILM" OR "CG

=> d que L42

L41 36 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON ("KARAOLIS D"/AU OR  
"KARAOLIS D K R"/AU OR "KARAOLIS DAVID"/AU OR "KARAOLIS DAVID  
K R"/AU)

L42 16 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L41 AND (GUAN? OR  
BIOFILM? OR BIO(W)FILM? OR "BIOFILM" OR "CGMP" OR "GMP")

=> d his L44

(FILE 'MEDLINE, BIOSIS, EMBASE, DRUGU' ENTERED AT 16:49:19 ON 19 OCT 2009)

L44 44 S L42

=> d que L44

L41 36 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON ("KARAOLIS D"/AU OR  
"KARAOLIS D K R"/AU OR "KARAOLIS DAVID"/AU OR "KARAOLIS DAVID  
K R"/AU)

L42 16 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L41 AND (GUAN? OR  
BIOFILM? OR BIO(W)FILM? OR "BIOFILM" OR "CGMP" OR "GMP")

L44 44 SEA L42

=> dup rem L42 L44

FILE 'HCAPLUS' ENTERED AT 17:02:30 ON 19 OCT 2009

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'MEDLINE' ENTERED AT 17:02:30 ON 19 OCT 2009

FILE 'BIOSIS' ENTERED AT 17:02:30 ON 19 OCT 2009

Copyright (c) 2009 The Thomson Corporation

FILE 'EMBASE' ENTERED AT 17:02:30 ON 19 OCT 2009

Copyright (c) 2009 Elsevier B.V. All rights reserved.

FILE 'DRUGU' ENTERED AT 17:02:30 ON 19 OCT 2009

COPYRIGHT (C) 2009 THOMSON REUTERS

PROCESSING COMPLETED FOR L42

PROCESSING COMPLETED FOR L44

L58 25 DUP REM L42 L44 (35 DUPLICATES REMOVED)

ANSWERS '1-16' FROM FILE HCAPLUS

ANSWERS '17-25' FROM FILE BIOSIS

**Inventor search results**

=&gt; d L58 1-25 ibib ab

L58 ANSWER 1 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 1

ACCESSION NUMBER: 2009:869974 HCAPLUS Full-textTITLE: c-di-**GMP** as a vaccine adjuvant enhances protection against systemic methicillin-resistant Staphylococcus aureus (MRSA) infectionAUTHOR(S): Hu, Dong-Liang; Narita, Kouji; Hyodo, Mamoru; Hayakawa, Yoshihiro; Nakane, Akio; **Karaolis, David K. R.**

CORPORATE SOURCE: Department of Microbiology and Immunology, Hirosaki University Graduate School of Medicine, Hirosaki, 036-8562, Japan

SOURCE: Vaccine (2009), 27(35), 4867-4873

CODEN: VACCDE; ISSN: 0264-410X

PUBLISHER: Elsevier Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Cyclic diguanylate (c-di-**GMP**) is a novel immunomodulator and immune enhancer that triggers a protective host innate immune response. The protective effect of c-di-**GMP** as a vaccine adjuvant against Staphylococcus aureus infection was investigated by s.c. (s.c.) vaccination with two different S. aureus antigens, clumping factor A (ClfA) and a nontoxic mutant staphylococcal enterotoxin C (mSEC), then i.v. (i.v.) challenge with viable methicillin-resistant S. aureus (MRSA) in a systemic infection model. Mice immunized with c-di-**GMP** plus mSEC or c-di-**GMP** plus ClfA vaccines then challenged with MRSA produced strong antigen-specific antibody responses demonstrating immunogenicity of the vaccines. Bacterial counts in the spleen and liver of c-di-**GMP** plus mSEC and c-di-**GMP** plus ClfA-immunized mice were significantly lower than those of control mice ( $P < 0.001$ ). Mice immunized with c-di-**GMP** plus mSEC or c-di-**GMP** plus ClfA showed significantly higher survival rates at day 7 (87.5%) than those of the non-immunized control mice (33.3%) ( $P < 0.05$ ). Furthermore, immunization of mice with c-di-**GMP** plus mSEC or c-di-**GMP** plus ClfA induced not only very high titers of IgG1 (IgG1), but c-di-**GMP** plus mSEC also induced significantly higher levels of IgG2a, IgG2b and IgG3 compared to alum adjuvant ( $P < 0.01$  and  $P < 0.001$ , resp.) and c-di-**GMP** plus ClfA induced significantly higher levels of IgG2a, IgG2b and IgG3 compared to alum adjuvant ( $P < 0.001$ ). Our results show that c-di-**GMP** should be developed as an adjuvant and immunotherapeutic to provide protection against systemic infection caused by S. aureus (MRSA).

L58 ANSWER 2 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 2

ACCESSION NUMBER: 2008:965252 HCAPLUS Full-text

DOCUMENT NUMBER: 150:349833

TITLE: c-di-**GMP** is an effective immunomodulator and vaccine adjuvant against pneumococcal infectionAUTHOR(S): Ogunniyi, Abiodun D.; Paton, James C.; Kirby, Alun C.; McCullers, Jonathan A.; Cook, Jan; Hyodo, Mamoru; Hayakawa, Yoshihiro; **Karaolis, David K. R.**

CORPORATE SOURCE: School of Molecular and Biomedical Science, University of Adelaide, 5005, Australia

SOURCE: Vaccine (2008), 26(36), 4676-4685

CODEN: VACCDE; ISSN: 0264-410X

PUBLISHER: Elsevier Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Cyclic diguanylate (c-di-GMP) is a unique bacterial intracellular signaling mol. capable of stimulating enhanced protective innate immunity against various bacterial infections. The effects of intranasal pretreatment with c-di-GMP, or i.p. coadministration of c-di-GMP with the pneumolysin toxoid (PdB) or pneumococcal surface protein A (PspA) before pneumococcal challenge, were investigated in mice. We found that c-di-GMP had no significant direct short-term effect on the growth rate of *Streptococcus pneumoniae* either in vitro or in vivo. However, intranasal pretreatment of mice with c-di-GMP resulted in a significant decrease in bacterial load in lungs and blood after serotypes 2 and 3 challenge, and a significant decrease in lung titers after serotype 4 challenge. Potential cellular mediators of these enhanced protective responses were identified in lungs and draining lymph nodes. I.p. coadministration of c-di-GMP with PdB or PspA before challenge resulted in significantly higher antigen-specific antibody titers and increased survival of mice, compared to that obtained with alum adjuvant. These findings demonstrate that local or systemic c-di-GMP administration stimulates innate and adaptive immunity against invasive pneumococcal disease. We propose that c-di-GMP can be used as an effective broad spectrum immunomodulator and vaccine adjuvant to prevent infectious diseases. OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD

(1 CITINGS)

REFERENCE COUNT: 50 THERE ARE 50 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L58 ANSWER 3 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 3

ACCESSION NUMBER: 2007:1122256 HCAPLUS Full-text

DOCUMENT NUMBER: 147:514687

TITLE: Cyclic di-GMP stimulates protective innate immunity in bacterial pneumonia

AUTHOR(S): Karaolis, David K. R.; Newstead, Michael W.; Zeng, Xianying; Hyodo, Mamoru; Hayakawa, Yoshihiro; Bhan, Urvashi; Liang, Hallie; Standiford, Theodore J.

CORPORATE SOURCE: Intragenics Research Institute, Havre de Grace, MD, 21078, USA

SOURCE: Infection and Immunity (2007), 75(10), 4942-4950  
CODEN: INFIBR; ISSN: 0019-9567

PUBLISHER: American Society for Microbiology

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Innate immunity is the primary mechanism by which extracellular bacterial pathogens are effectively cleared from the lung. We have previously shown that cyclic di-GMP (c-di-GMP [c-diguanylate]) is a novel small mol. immunomodulator and immunostimulatory agent that triggers protective host innate immune responses. Using a murine model of bacterial pneumonia, we show that local intranasal (i.n.) or systemic s.c. administration of c-di-GMP prior to intratracheal (i.t.) challenge with *Klebsiella pneumoniae* stimulates protective immunity against infection. Specifically, i.n. or s.c. administration of c-di-GMP 48 and 24 h prior to i.t. *K. pneumoniae* challenge resulted in significantly increased survival. Pretreatment with c-di-GMP resulted in a 5-fold reduction in bacterial CFU in the lung ( $P < 0.05$ ) and an impressive  $> 1000$ -fold decrease in CFU in the blood ( $P < 0.01$ ). C-di-GMP administration stimulated a robust innate response to bacterial challenge, characterized by enhanced accumulation of neutrophils and  $\alpha\beta$  T cells, as well as activated NK and  $\alpha\beta$  T lymphocytes, which was associated with earlier and more vigorous expression of chemokines and type I cytokines. Moreover, lung macrophages recovered from *Klebsiella*-infected mice pretreated with c-di-GMP expressed greater quantities of inducible nitric oxide synthase and nitric oxide ex vivo than did macrophages isolated from infected mice pretreated with the control, c-GMP. These findings demonstrate that c-di-GMP delivered in either a compartmentalized or systemic fashion stimulates protective innate immunity in the lung and protects mice against bacterial invasion. We propose that the cyclic dinucleotide c-di-GMP may be used clin. as an effective immunomodulator, immune enhancer, and vaccine

adjuvant to protect against respiratory infection and pneumonia in humans and animals. OS.CITING REF COUNT: 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD

(9 CITINGS)

REFERENCE COUNT: 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L58 ANSWER 4 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 4

ACCESSION NUMBER: 2007:124392 HCAPLUS Full-text

DOCUMENT NUMBER: 146:204267

TITLE: Bacterial c-di-GMP Is an Immunostimulatory Molecule

AUTHOR(S): **Karaolis, David K. R.**; Means, Terry K.; Yang, De; Takahashi, Munehisu; Yoshimura, Teizo; Muraille, Eric; Philpott, Dana; Schroeder, John T.; Hyodo, Mamoru; Hayakawa, Yoshihiro; Talbot, Brian G.; Brouillette, Eric; Malouin, Francois

CORPORATE SOURCE: Intragenics Research Institute, Havre de Grace, MD, 21078, USA

SOURCE: Journal of Immunology (2007), 178(4), 2171-2181  
CODEN: JOIMA3; ISSN: 0022-1767

PUBLISHER: American Association of Immunologists

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Cyclic diguanylate (c-di-GMP) is a bacterial intracellular signaling mol. The authors have shown that treatment with exogenous c-di-GMP inhibits Staphylococcus aureus infection in a mouse model. The authors now report that c-di-GMP is an immunomodulator and immunostimulatory mol. Intramammary treatment of mice with c-di-GMP 12 and 6 h before S. aureus challenge gave a protective effect and a 10,000-fold reduction in CFUs in tissues. I.m. vaccination of mice with c-di-GMP coinjected with S. aureus clumping factor A (ClfA) Ag produced serum with significantly higher anti-ClfA IgG Ab titers compared with ClfA alone. I.p. injection of mice with c-di-GMP activated monocyte and granulocyte recruitment. Human immature dendritic cells (DCs) cultured in the presence of c-di-GMP showed increased expression of costimulatory mols. CD80/CD86 and maturation marker CD83, increased MHC class II and cytokines and chemokines such as IL-12, IFN- $\gamma$ , IL-8, MCP-1, IFN- $\gamma$ -inducible protein 10, and RANTES, and altered expression of chemokine receptors including CCR1, CCR7, and CXCR4. C-di-GMP-matured DCs demonstrated enhanced T cell stimulatory activity. C-di-GMP activated p38 MAPK in human DCs and ERK phosphorylation in human macrophages. C-di-GMP is stable in human serum. The authors propose that cyclic dinucleotides like c-di-GMP can be used clin. in humans and animals as an immunomodulator, immune enhancer, immunotherapeutic, immunoprophylactic, or vaccine adjuvant.

OS.CITING REF COUNT: 12 THERE ARE 12 CAPLUS RECORDS THAT CITE THIS RECORD (12 CITINGS)

REFERENCE COUNT: 58 THERE ARE 58 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L58 ANSWER 5 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 5

ACCESSION NUMBER: 2005:714247 HCAPLUS Full-text

DOCUMENT NUMBER: 143:205833

TITLE: 3',5'-Cyclic diguanylic acid reduces the virulence of biofilm-forming Staphylococcus aureus strains in a mouse model of mastitis infection

AUTHOR(S): Brouillette, Eric; Hyodo, Mamoru; Hayakawa, Yoshihiro; **Karaolis, David K. R.**; Malouin, Francois

CORPORATE SOURCE: Centre d'Etude et de Valorisation de la Diversite Microbienne (CEVDM), Departement de biologie, Faculte des sciences, Universite de Sherbrooke, Sherbrooke, QC, J1K 2R1, Can.

SOURCE: Antimicrobial Agents and Chemotherapy (2005), 49(8),  
3109-3113  
CODEN: AMACCQ; ISSN: 0066-4804  
PUBLISHER: American Society for Microbiology  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB The cyclic dinucleotide 3',5'-cyclic diguanylic acid (c-di-GMP) is a naturally occurring small mol. that regulates important signaling systems in bacteria. The authors have recently shown that c-di-GMP inhibits *Staphylococcus aureus* **biofilm** formation in vitro and its adherence to HeLa cells. The authors now report that c-di-GMP treatment has an antimicrobial and antipathogenic activity in vivo and reduces, in a dose-dependent manner, bacterial colonization by **biofilm**-forming *S. aureus* strains in a mouse model of mastitis infection. Intramammary injections of 5 and 50 nmol of c-di-GMP decreased colonization (bacterial CFU) per g of gland by 0.79 (P > 0.05) and 1.44 (P < 0.01) logs, resp., whereas 200-nmol doses allowed clearance of the bacteria below the detection limit with a reduction of more than 4 logs (P < 0.001) compared to the untreated control groups. These results indicate that cyclic dinucleotides potentially represent an attractive and novel drug platform which could be used alone or in combination with other agents or drugs in the prevention, treatment, or control of infection. OS.CITING REF

COUNT: 24 THERE ARE 24 CAPLUS RECORDS THAT CITE THIS

RECORD (24 CITINGS)

REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L58 ANSWER 6 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 6

ACCESSION NUMBER: 2005:229578 HCAPLUS Full-text

DOCUMENT NUMBER: 142:426617

TITLE: c-di-GMP (3'-5'-cyclic diguanylic acid)  
inhibits *Staphylococcus aureus* cell-cell interactions  
and **biofilm** formation

AUTHOR(S): Karaolis, David K. R.; Rashid, Mohammed H.;  
Chythanya, Rajanna; Luo, Wensheng; Hyodo, Mamoru;  
Hayakawa, Yoshihiro

CORPORATE SOURCE: Department of Epidemiology and Preventive Medicine,  
University of Maryland School of Medicine, Baltimore,  
MD, 21201, USA

SOURCE: Antimicrobial Agents and Chemotherapy (2005), 49(3),  
1029-1038

CODEN: AMACCQ; ISSN: 0066-4804

PUBLISHER: American Society for Microbiology

DOCUMENT TYPE: Journal

LANGUAGE: English

AB *Staphylococcus aureus* is an important pathogen of humans and animals, and antibiotic resistance is a public health concern. **Biofilm** formation is essential in virulence and pathogenesis, and the ability to resist antibiotic treatment results in difficult-to-treat and persistent infections. As such, novel antimicrobial approaches are of great interest to the scientific, medical, and agriculture communities. We recently proposed that modulating levels of the cyclic dinucleotide signaling mol., c-di-GMP (cyclic diguanylate [3',5'-cyclic diguanylic acid], cGpGp), has utility in regulating phenotypes of prokaryotes. We report that extracellular c-di-GMP shows activity against human clin. and bovine intramammary mastitis isolates of *S. aureus*, including methicillin-resistant *S. aureus* (MRSA) isolates. We show that chemical synthesized c-di-GMP is soluble and stable in water and physiol. saline and stable following boiling and exposure to acid and alkali. Treatment of *S. aureus* with extracellular c-di-GMP inhibited cell-to-cell (intercellular) adhesive interactions in liquid medium and reduced (>50%) **biofilm** formation in human and bovine isolates compared to untreated controls. C-di-GMP inhibited the adherence of *S. aureus* to human epithelial HeLa cells. The cyclic nucleotide analogs cGMP and cAMP had a lesser inhibitory effect on **biofilms**, while

5'-**GMP** had no major effect. We propose that cyclic dinucleotides such as c-di-**GMP**, used either alone or in combination with other antimicrobial agents, represent a novel and attractive approach in the development of intervention strategies for the prevention of **biofilms** and the control and treatment of infection.

OS.CITING REF COUNT: 30 THERE ARE 30 CAPLUS RECORDS THAT CITE THIS RECORD (30 CITINGS)  
 REFERENCE COUNT: 64 THERE ARE 64 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L58 ANSWER 7 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 7  
 ACCESSION NUMBER: 2007:662269 HCAPLUS Full-text  
 DOCUMENT NUMBER: 148:496266  
 TITLE: Chemical behavior of bis(3'-5')diguanylic acid in aqueous solutions  
 AUTHOR(S): Hyodo, Mamoru; Sato, Yumi; Hayakawa, Yoshihiro; **Karaolis, David K. R.**  
 CORPORATE SOURCE: Graduate School of Information Science/Human Informatics, Nagoya University, Chikusa, Nagoya, 464-8601, Japan  
 SOURCE: Nucleic Acids Symposium Series (2005), (49), 117-118  
 CODEN: NASSCJ  
 URL: <http://nass.oxfordjournals.org/content/vol49/issue1/index.dtl>  
 PUBLISHER: Oxford University Press  
 DOCUMENT TYPE: Journal; (online computer file)  
 LANGUAGE: English

AB A unique behavior of bis(3'-5')diguanylic acid (c-di-**GMP**) under some conditions is described. It exists as the monomer in aprotic organic solvents such as DMSO. By contrast, it smoothly aggregates in water and in low-concentration aqueous solns. of some salts, such as sodium chloride and ammonium acetate, to give a mixture of many aggregates. The resulting multiple aggregates converge to the single compound (provably the monomer) in a >154 mM (0.9%) sodium chloride aqueous solution, in a >100 mM ammonium acetate buffer, and in a >100 mM phosphate buffer.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
 REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L58 ANSWER 8 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 8  
 ACCESSION NUMBER: 2005:150086 HCAPLUS Full-text  
 DOCUMENT NUMBER: 142:329238  
 TITLE: 3',5'-Cyclic diguanylic acid (c-di-**GMP**) inhibits basal and growth factor-stimulated human colon cancer cell proliferation  
 AUTHOR(S): **Karaolis, David K. R.**; Cheng, Kunrong; Lipsky, Michael; Elnabawi, Ahmed; Catalano, Jennifer; Hyodo, Mamoru; Hayakawa, Yoshihiro; Raufman, Jean-Pierre  
 CORPORATE SOURCE: Department of Epidemiology and Preventive Medicine, University of Maryland School of Medicine, Baltimore, MD, 21201, USA  
 SOURCE: Biochemical and Biophysical Research Communications (2005), 329(1), 40-45  
 CODEN: BBRCA9; ISSN: 0006-291X  
 PUBLISHER: Elsevier  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB The novel cyclic dinucleotide, 3',5'-cyclic diguanylic acid, cGpGp (c-di-**GMP**), is a naturally occurring small mol. that regulates important signaling mechanisms in prokaryotes. Recently, we showed that c-di- **GMP** has "drug-like"

properties and that c-di-GMP treatment might be a useful antimicrobial approach to attenuate the virulence and pathogenesis of *Staphylococcus aureus* and prevent or treat infection. In the present communication, we report that c-di-GMP ( $\geq 50 \mu\text{M}$ ) has striking properties regarding inhibition of cancer cell proliferation in vitro. c-di-GMP inhibits both basal and growth factor (acetylcholine and epidermal growth factor)-induced cell proliferation of human colon cancer (H508) cells. Toxicity studies revealed that exposure of normal rat kidney cells and human neuroblastoma cells to c-di-GMP at biol. relevant doses showed no lethal cytotoxicity. Cyclic dinucleotides, such as c-di-GMP, represent an attractive and novel "drug-platform technol." that can be used not only to develop new antimicrobial agents, but also to develop novel therapeutic agents to prevent or treat cancer.

OS.CITING REF COUNT: 18 THERE ARE 18 CAPLUS RECORDS THAT CITE THIS RECORD (18 CITINGS)  
 REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L58 ANSWER 9 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 9

ACCESSION NUMBER: 2004:50869 HCAPLUS Full-text

DOCUMENT NUMBER: 140:178223

TITLE: Role of exopolysaccharide, the rugose phenotype and VpsR in the pathogenesis of epidemic *Vibrio cholerae*  
 AUTHOR(S): Rashid, Mohammed H.; Rajanna, Chythanya; Zhang, Dalin; Pasquale, Vincenzo; Magder, Laurence S.; Ali, Afsar; Dumontet, Stefano; **Karaolis, David K. R.**

CORPORATE SOURCE: Department of Epidemiology and Preventive Medicine, University of Maryland School of Medicine, Baltimore, MD, 21201, USA

SOURCE: FEMS Microbiology Letters (2004), 230(1), 105-113  
 CODEN: FMLED7; ISSN: 0378-1097

PUBLISHER: Elsevier Science B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB *Vibrio cholerae*, the causative agent of cholera, can produce an exopolysaccharide (EPS). Some strains can also phenotypically switch from a smooth to a 'rugose' phenotype characterized by small wrinkled colonies, overprodn. of EPS, increased **biofilm** formation in vitro and increased resistance to various stressful conditions. High frequency switching to the rugose phenotype is more common in epidemic strains than in non-pathogenic strains, suggesting EPS production and the rugose phenotype are important in cholera epidemiol. VpsR up-regulates *Vibrio* polysaccharide (VPS) genes and the synthesis of extracellular EPS (VPS). However, the function of VPS, the rugose phenotype and VpsR in pathogenesis is not well understood. The authors report that rugose strains of both classical and El Tor biotypes of epidemic *V. cholerae* are defective in the in vitro production of extracellular collagenase activity. In vivo studies in rabbit ileal loops suggest that VpsR mutants are attenuated in reactogenicity. Intestinal colonization studies in infant mice suggest that VPS production, the rugose phenotype and VpsR have a role in pathogenesis. The results indicate that regulated VPS production is important for promoting in vivo **biofilm** formation and pathogenesis. Addnl., VpsR might regulate genes with roles in virulence. Rugose strains appear to be a subpopulation of cells that might act as a 'helper' phenotype promoting the pathogenesis of certain strains. These studies provide new insight into the potential role of VPS, the rugose phenotype and VpsR in the pathogenesis of epidemic *V. cholerae*. OS.CITING REF COUNT: 8 THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD

(8 CITINGS)  
 REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L58 ANSWER 10 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 10

ACCESSION NUMBER: 2003:688110 HCAPLUS Full-text

DOCUMENT NUMBER: 140:14616  
 TITLE: Analysis of the *Vibrio* pathogenicity island-encoded Mop protein suggests a pleiotropic role in the virulence of epidemic *Vibrio cholerae*  
 AUTHOR(S): Zhang, Dalin; Rajanna, Chythanya; Sun, Weiyun; **Karaolis, David K. R.**  
 CORPORATE SOURCE: Department of Epidemiology and Preventive Medicine, University of Maryland School of Medicine, Baltimore, MD, 21201, USA  
 SOURCE: FEMS Microbiology Letters (2003), 225(2), 311-318  
 CODEN: FMLED7; ISSN: 0378-1097  
 PUBLISHER: Elsevier Science B.V.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Epidemic *Vibrio cholerae* contain a large essential virulence gene cluster called the *Vibrio* pathogenicity island (VPI). We recently reported that no in vitro difference in virulence was found in El Tor strain N16961 containing a mutation in the VPI-encoded mop gene but this mutant was hypervirulent and reactogenic in rabbit ileal loops. In this paper, we report in vitro studies showing that independent Mop mutants of strain 3083 are significantly attenuated (.apprx.40-fold) in cholera toxin (CT) production and have significantly increased motility and **biofilm** forming ability but appear to be unaffected in TcpA, hemagglutinin protease and hemolysin compared to their parent. The 3083 Mop mutant showed a 100-fold decrease in its in vivo intestinal colonization ability in the infant mouse competition assays. While reverse transcription polymerase chain reaction and phenotypic studies of a mop plasmid in both mutant and wild-type backgrounds suggest Mop is expressed by the plasmid, the differences in CT and **biofilm** formation could not be restored in any of the mutants. The inability to complement the Mop mutants in trans may be due either to the selection of secondary mutations or to mop possibly being part of an operon. Our findings that Mop is associated with CT, motility, **biofilm** formation and intestinal colonization support a hypothesis in which Mop has a pleiotropic role in the pathogenesis and persistence of epidemic *V. cholerae*. OS.CITING REF COUNT: 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD

(4 CITINGS)

REFERENCE COUNT: 48 THERE ARE 48 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L58 ANSWER 11 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 11

ACCESSION NUMBER: 2003:808644 HCAPLUS Full-text  
 DOCUMENT NUMBER: 140:54341  
 TITLE: Identification of genes involved in the switch between the smooth and rugose phenotypes of *Vibrio cholerae*  
 AUTHOR(S): Rashid, Mohammed H.; Rajanna, Chythanya; Ali, Afsar; **Karaolis, David K. R.**  
 CORPORATE SOURCE: Department of Epidemiology and Preventive Medicine, University of Maryland School of Medicine, Baltimore, MD, 21201, USA  
 SOURCE: FEMS Microbiology Letters (2003), 227(1), 113-119  
 CODEN: FMLED7; ISSN: 0378-1097  
 PUBLISHER: Elsevier Science B.V.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB *Vibrio cholerae* can switch to a rugose' phenotype characterized by an exopolysaccharide (EPS) matrix, wrinkled colony morphol., increased **biofilm** formation and increased survival under specific conditions. The vps gene cluster responsible for the biosynthesis of the rugose EPS (rEPS) is pos. regulated by VpsR. We recently identified media (APW#3) promoting EPS production and the rugose phenotype and found epidemic strains switch at a higher frequency than non-pathogenic strains, suggesting this switch and the rugose phenotype are important



in cholera epidemiol. In this study, transposon mutagenesis on a smooth V. cholerae strain was used to identify mutants that were unable to shift to the rugose phenotype under inducing conditions to better understand the mol. basis of the switch. We identified vpsR, gale and vps previously associated with the rugose phenotype, and also identified genes not previously associated with the phenotype, including rfbD and rfbE having roles in LPS (lipopolysaccharide) synthesis and aroB and aroK with roles in aromatic amino acid synthesis. Addnl., a mutation in amiB encoding N-acetylmuramoyl-L-alanine amidase caused defects in the switch, motility and cell morphol. We also found that a gene encoding a novel regulatory protein we termed RocS (regulation of cell signaling) containing a GGDEF and EAL domains and associated with c-di-GMP levels is important for the rugose phenotype, EPS, **biofilm** formation and motility. We propose that modulation of cyclic dinucleotide (e.g. c-di-GMP) levels might have application in regulating various phenotypes of prokaryotes. Our study shows the mol. complexity of the switch between the smooth and rugose phenotypes of V. cholerae and may be relevant to similar phenotypes in other species. OS.CITING REF COUNT: 51 THERE ARE 51 CAPLUS RECORDS THAT CITE THIS

REFERENCE COUNT: 55 RECORD (52 CITINGS)  
THERE ARE 55 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L58 ANSWER 12 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 12

ACCESSION NUMBER: 2002:862053 HCAPLUS Full-text

DOCUMENT NUMBER: 138:119678

TITLE: High-frequency rugose exopolysaccharide production by Vibrio cholerae

AUTHOR(S): Ali, Afsar; Rashid, Mohammed H.; **Karaolis, David K. R.**

CORPORATE SOURCE: Department of Epidemiology and Preventive Medicine, University of Maryland School of Medicine, Baltimore, MD, 21201, USA

SOURCE: Applied and Environmental Microbiology (2002), 68(11), 5773-5778

CODEN: AEMIDF; ISSN: 0099-2240

PUBLISHER: American Society for Microbiology

DOCUMENT TYPE: Journal

LANGUAGE: English

AB V. cholerae can shift to a "rugose" phenotype, thereby producing copious exopolysaccharide (EPS), which promotes its environmental survival and persistence. We report conditions that promote high-frequency rugose EPS production (HFRP), whereby cells switch at high frequency ( $\leq 80\%$ ) to rugose EPS production. HFRP appeared to be more common in clin. strains, as HFRP was found in 6 of 19 clin. strains (32%) (including classical, El Tor, and non-O1 strains) but in only 1 of 16 environmental strains (6%). Differences were found between strains in rugose colony morphol., conditions promoting HFRP, the frequency of rugose-to-smooth (R-S) cell reversion, and **biofilm** formation. We propose that rugose EPS and HFRP provide an evolutionary and adaptive advantage to specific epidemic V. cholerae strains for increased persistence in the environment. OS.CITING REF COUNT: 25 THERE ARE 25 CAPLUS RECORDS THAT CITE THIS

REFERENCE COUNT: 30 RECORD (25 CITINGS)  
THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L58 ANSWER 13 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2007:1396598 HCAPLUS Full-text

DOCUMENT NUMBER: 148:24432

TITLE: Method for stimulating the immune, inflammatory or neuroprotective response

INVENTOR(S): **Karaolis, David K. R.**

PATENT ASSIGNEE(S): USA

10/565,591

SOURCE: U.S. Pat. Appl. Publ., 60pp., Cont.-in-part of U.S. Ser. No. 79,886.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20070281897	A1	20071206	US 2007-669006	20070130
US 7592326	B2	20090922		
AU 2005221717	A1	20050922	AU 2005-221717	20050315
CA 2559802	A1	20050922	CA 2005-2559802	20050315
AU 2005222626	A1	20050929	AU 2005-222626	20050315
CA 2560058	A1	20050929	CA 2005-2560058	20050315
US 20060040887	A1	20060223	US 2005-79886	20050315
US 7569555	B2	20090804		
EP 1729781	A1	20061213	EP 2005-727318	20050315
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR				
EP 1740192	A2	20070110	EP 2005-753223	20050315
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR				
JP 2007529531	T	20071025	JP 2007-503996	20050315
JP 2007529532	T	20071025	JP 2007-503997	20050315
PRIORITY APPLN. INFO.:				
			US 2004-552721P	P 20040315
			US 2004-563692P	P 20040420
			US 2005-79886	A2 20050315
			WO 2005-US8447	W 20050315
			WO 2005-US8448	W 20050315

AB Cyclic di-**GMP**, or a cyclic dinucleotide analog thereof that has the same effect as cyclic di-**GMP**, stimulates or enhances immune or inflammatory response in a patient or enhances the immune response to a vaccine by serving as an adjuvant. Cyclic di-**GMP**, or a cyclic dinucleotide analog thereof, also has neuroprotective properties for use as a neuroprotective agent to inhibit, treat, or ameliorate the effects of injuries, diseases,.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L58 ANSWER 14 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2006:351606 HCAPLUS Full-text

DOCUMENT NUMBER: 145:189078

TITLE: Organic synthesis, chemical properties, and biological activities of cyclic bis(3'-5')diguanylic acid (c-di-**GMP**) and its analogs

AUTHOR(S): Hyodo, Mamoru; Hayakawa, Yoshihiro; **Karaolis, David K. R.**

CORPORATE SOURCE: Graduate School of Human Informatics/Information Science, CREST/JST, Nagoya University, Chikusa, Nagoya, 464-8601, Japan

SOURCE: Yuki Gosei Kagaku Kyokaishi (2006), 64(4), 359-370  
 CODEN: YGKKAE; ISSN: 0037-9980

PUBLISHER: Yuki Gosei Kagaku Kyokai

DOCUMENT TYPE: Journal; General Review

LANGUAGE: Japanese

AB A review. This paper describes efficient synthesis, chemical behaviors, and biol. activities of cyclic bis(3'-5')diguanylic acid (c-di-**GMP**) and its analogs, including cyclic bis(3'-5')**guanylic**-inosinic acid (c-GpIp), cyclic bis(3'-5')**guanylic**-adenylic acid (c-GpAp), and bis(3'-5')diguanylic acid

monophosphorothioate (c-GpGs). C-di- **GMP** was synthesized via two methods. Between the two methods, one method is more effective, particularly, for large-scale (gram-scale) synthesis to obtain the target compound in a high yield. While, c-GpIp, c-GpAp, and c-GpGs were synthesized via similar strategies. Studies on chemical behaviors of c-di-**GMP** indicated that these cyclic dinucleotides exist as the monomers in aprotic solvents such as DMSO. By contrast, it was shown that c-di-**GMP** smoothly aggregates to form a mixture of many compds. in water, in < 0.9% sodium chloride solns., in < 100 mM phosphate buffer solns., and in < 100 mM ammonium acetate buffer solns. All aggregated compds. smoothly revert to a single compound (probably an aggregate) by dissolving in a 0.9% sodium chloride solution (a physiol. salt solution), a > 100 mM phosphate buffer solution, or a > 100 mM ammonium acetate buffer solution. Biol. investigation disclosed some novel activities of c-di-**GMP**, such as inhibition of **biofilm** formation of *Staphylococcus aureus*, inhibition of basal and growth factor stimulated human colon cancer cell proliferation, and reduction of the villus of **biofilm**-formed *Staphylococcus aureus* in a mouse model. OS.CITING REF COUNT: 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD

(4 CITINGS)

L58 ANSWER 15 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2005:300236 HCAPLUS Full-text

DOCUMENT NUMBER: 142:367640

TITLE: Method for attenuating virulence of microbial pathogens and inhibiting microbial **biofilm** formation by using c-di-**GMP** and cyclic dinucleotide analogs

INVENTOR(S): **Karaolis, David K. R.**

PATENT ASSIGNEE(S): University of Maryland, USA

SOURCE: PCT Int. Appl., 118 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005030186	A2	20050407	WO 2004-US23498	20040722
WO 2005030186	A3	20050714		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2004275696	A1	20050407	AU 2004-275696	20040722
CA 2533873	A1	20050407	CA 2004-2533873	20040722
EP 1651242	A2	20060503	EP 2004-809506	20040722
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK			
JP 2007500697	T	20070118	JP 2006-521912	20040722
US 20070244059	A1	20071018	US 2006- <b>565591</b>	20061006
PRIORITY APPLN. INFO.:			US 2003-490029P	P 20030728
			WO 2004-US23498	W 20040722

AB The present invention relates to the use of the cyclic dinucleotide c-di- **GMP**

and cyclic dinucleotide analogs thereof in a method for attenuating virulence of a microbial pathogen or for inhibiting or reducing colonization by a microbial pathogen. This method further inhibits microbial **biofilm** formation and is capable of treating bacterial infections. The microbial colonization or **biofilm** formation inhibited or reduced may be on the skin or on nasal or mucosal surface. The microbial colonization or **biofilm** formation inhibited can also be on the surfaces of medical devices, especially those in close contact with the patient, as well on the surfaces of industrial and construction material where microbial colonization and **biofilm** formation is of concern.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)

L58 ANSWER 16 OF 25 HCAPLUS COPYRIGHT 2009 ACS on STN  
ACCESSION NUMBER: 2005:1005971 HCAPLUS Full-text  
DOCUMENT NUMBER: 143:279369  
TITLE: Method using cyclic di-GMP or cyclic  
dinucleotide analog thereof for inhibiting cancer cell  
proliferation or increasing cancer cell apoptosis  
INVENTOR(S): Karaolis, David K.R.; Raufman, Jean-Pierre  
PATENT ASSIGNEE(S): USA  
SOURCE: U.S. Pat. Appl. Publ., 22 pp.  
CODEN: USXXCO  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20050203051	A1	20050915	US 2005-79779	20050315
AU 2005221717	A1	20050922	AU 2005-221717	20050315
CA 2559802	A1	20050922	CA 2005-2559802	20050315
WO 2005087238	A2	20050922	WO 2005-US8447	20050315
WO 2005087238	A3	20060309		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2005222626	A1	20050929	AU 2005-222626	20050315
CA 2560058	A1	20050929	CA 2005-2560058	20050315
WO 2005089777	A1	20050929	WO 2005-US8448	20050315
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1729781	A1	20061213	EP 2005-727318	20050315

10/565,591

R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,  
IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR  
EP 1740192 A2 20070110 EP 2005-753223 20050315  
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,  
IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR  
JP 2007529531 T 20071025 JP 2007-503996 20050315  
JP 2007529532 T 20071025 JP 2007-503997 20050315  
PRIORITY APPLN. INFO.: US 2004-552721P P 20040315  
US 2004-563692P P 20040420  
WO 2005-US8447 W 20050315  
WO 2005-US8448 W 20050315

AB Cyclic di-**GMP** or cyclic dinucleotide analogs thereof can be used to inhibit cancer cell proliferation or to increase cancer cell apoptosis in vitro as well as in vivo in a patient. OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD

(2 CITINGS)

L58 ANSWER 17 OF 25 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on STN

ACCESSION NUMBER: 2009:518645 BIOSIS Full-text  
DOCUMENT NUMBER: PREV200900519748  
TITLE: Method for stimulating the immune, inflammatory or neuroprotective response.  
AUTHOR(S): **Karaolis, David K. R.** [Inventor]; Anonymous  
CORPORATE SOURCE: Baltimore, MD 21210 USA  
PATENT INFORMATION: US 07569555 20090804  
SOURCE: Official Gazette of the United States Patent and Trademark Office Patents, (AUG 4 2009)  
CODEN: OGUPE7. ISSN: 0098-1133.  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
ENTRY DATE: Entered STN: 2 Sep 2009  
Last Updated on STN: 2 Sep 2009

AB Cycic di-**GMP**, or a cyclic dinucleotide analogue thereof that has the same effect as cyclic di-**GMP**, stimulates or enhances immune or inflammatory response in a patient or enhances the immune response to a vaccine by serving as an adjuvant. Cyclic di-**GMP**, or a cyclic dinucleotide analogue thereof, also has neuroprotective properties for use as a neuroprotective agent to inhibit, treat, or ameliorate the effects of injuries, diseases, disorders or conditions that result in neurodegeneration.

L58 ANSWER 18 OF 25 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on STN

ACCESSION NUMBER: 2008:193052 BIOSIS Full-text  
DOCUMENT NUMBER: PREV200800188893  
TITLE: c-di-**GMP** stimulates protective innate immunity in bacterial pneumonia.  
AUTHOR(S): **Karaolis, D. K. R.** [Reprint Author]; Newstead, M. W.; Zeng, X.; Liang, H.; Hyodo, M.; Hayakawa, Y.; Standiford, T. J.  
CORPORATE SOURCE: Intragen Res Inst, Havre De Grace, MD USA  
SOURCE: Abstracts of the General Meeting of the American Society for Microbiology, (2007) Vol. 107, pp. 266.  
Meeting Info.: 107th General Meeting of the American-Society-for-Microbiology. Toronto, CANADA. 2007,. Amer Soc Microbiol.  
ISSN: 1060-2011.  
DOCUMENT TYPE: Conference; (Meeting)  
Conference; (Meeting Poster)

10/565,591

LANGUAGE: English  
ENTRY DATE: Entered STN: 19 Mar 2008  
Last Updated on STN: 19 Mar 2008

L58 ANSWER 19 OF 25 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on  
STN

ACCESSION NUMBER: 2008:237637 BIOSIS Full-text  
DOCUMENT NUMBER: PREV200800233680  
TITLE: c-di-**GMP** is an immunostimulatory molecule with  
prophylactic and adjuvant activity.  
AUTHOR(S): **Karaolis, D. K. R.** [Reprint Author]; Means, T.  
K.; Brouillette, E.; Talbot, B. G.; Yang, D.; Muraille, E.;  
Hyodo, M.; Hayakawa, Y.; Malouin, F.  
CORPORATE SOURCE: Univ Maryland, Baltimore, MD 21201 USA  
SOURCE: Abstracts of the General Meeting of the American Society  
for Microbiology, (2006) Vol. 106, pp. 235.  
Meeting Info.: 106th General Meeting of the  
American-Society-for-Microbiology. Orlando, FL, USA. May 21  
-25, 2006. Amer Soc Microbiol.  
ISSN: 1060-2011.  
DOCUMENT TYPE: Conference; (Meeting)  
Conference; Abstract; (Meeting Abstract)  
LANGUAGE: English  
ENTRY DATE: Entered STN: 2 Apr 2008  
Last Updated on STN: 2 Apr 2008

L58 ANSWER 20 OF 25 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on  
STN

ACCESSION NUMBER: 2005:415877 BIOSIS Full-text  
DOCUMENT NUMBER: PREV200510201468  
TITLE: c-di-**GMP** as a novel anti-**biofilm** agent  
against Staphylococcus aureus.  
AUTHOR(S): **Karaolis, D. K. R.** [Reprint Author]; Rashid, M.  
H.; Rajanna, C.; Buckles, E.; Luo, W.; Hyodo, M.; Hayakawa,  
Y.  
SOURCE: Abstracts of the Interscience Conference on Antimicrobial  
Agents and Chemotherapy, (OCT-NOV 2004) Vol. 44, pp. 203.  
Meeting Info.: 44th Interscience Conference on  
Antimicrobial Agents and Chemotherapy. Washington, DC, USA.  
October 30 -November 02, 2004.  
ISSN: 0733-6373.  
DOCUMENT TYPE: Conference; (Meeting)  
Conference; (Meeting Poster)  
LANGUAGE: English  
ENTRY DATE: Entered STN: 19 Oct 2005  
Last Updated on STN: 19 Oct 2005

L58 ANSWER 21 OF 25 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on  
STN

ACCESSION NUMBER: 2007:335423 BIOSIS Full-text  
DOCUMENT NUMBER: PREV200700323424  
TITLE: Regulation of Vibrio cholerae **biofilm** formation  
and intestinal colonization by Vibrio pathogenicity island  
recombinases.  
AUTHOR(S): Rajanna, C. [Reprint Author]; Rashid, M. H.; **Karaolis,**  
**D. K. R.**  
CORPORATE SOURCE: Univ Maryland, Sch Med, Baltimore, MD 21201 USA  
SOURCE: Abstracts of the General Meeting of the American Society  
for Microbiology, (2004) Vol. 104, pp. 103.  
Meeting Info.: 104th General Meeting of the

American-Society-for-Microbiology. New Orleans, LA, USA.  
 May 23 -27, 2004. Amer Soc Microbiol.  
 ISSN: 1060-2011.

DOCUMENT TYPE: Conference; (Meeting)  
 Conference; Abstract; (Meeting Abstract)  
 LANGUAGE: English  
 ENTRY DATE: Entered STN: 30 May 2007  
 Last Updated on STN: 30 May 2007

L58 ANSWER 22 OF 25 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on  
 STN

ACCESSION NUMBER: 2003:544375 BIOSIS Full-text  
 DOCUMENT NUMBER: PREV200300546093  
 TITLE: The VPI-encoded int and VpiT of epidemic Vibrio cholerae  
 have roles in high frequency rugose exopolysaccharide  
 production (HFRP).  
 AUTHOR(S): Rajanna, C. [Reprint Author]; **Karaolis, D. K. R.**  
 [Reprint Author]  
 CORPORATE SOURCE: University of Maryland, Baltimore, MD, USA  
 SOURCE: Abstracts of the General Meeting of the American Society  
 for Microbiology, (2003) Vol. 103, pp. J-016.  
<http://www.asmsa.org/mtgsrc/generalmeeting.htm>. cd-rom.  
 Meeting Info.: 103rd American Society for Microbiology  
 General Meeting. Washington, DC, USA. May 18-22, 2003.  
 American Society for Microbiology.  
 ISSN: 1060-2011 (ISSN print).

DOCUMENT TYPE: Conference; (Meeting)  
 Conference; Abstract; (Meeting Abstract)  
 LANGUAGE: English  
 ENTRY DATE: Entered STN: 19 Nov 2003  
 Last Updated on STN: 19 Nov 2003

AB Vibrio cholerae causes the epidemic diarrheal disease called cholera. V.  
 cholerae can shift to a rugose colony phenotype in which the cells produce  
 copious amounts of exopolysaccharide (EPS). The production of this EPS  
 promotes the persistence of V. cholerae as it promotes **biofilm** formation and  
 increased resistance to various stresses such as acid, chlorine, UV light, and  
 complement. We have shown that some strains of V. cholerae can shift at high  
 frequency to produce EPS in a process we call high frequency rugose EPS  
 production (HFRP). HFRP was more common in clinical V. cholerae strains than  
 in nonpathogenic strains suggesting that EPS production and HFRP is important  
 in epidemic strains and might provide these strains with an adaptive advantage  
 in certain niches. Epidemic strains of V. cholerae contain the Vibrio  
 pathogenicity island (VPI) which is usually absent from nonpathogenic strains  
 and which is essential for virulence. We show that the VPI-encoded integrase  
 (int) and transposase-like (vpiT) genes affect EPS production and have roles  
 in the switch to HFRP. A VpiT mutant in particular is significantly reduced  
 in its ability to undergo HFRP and this defect can be complemented by a  
 plasmid containing vpiT. Since genes on the VPI can affect HFRP, our results  
 suggest that the VPI confers both virulence and increased environmental  
 persistence in epidemic V. cholerae strains.

L58 ANSWER 23 OF 25 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on  
 STN

ACCESSION NUMBER: 2003:544373 BIOSIS Full-text  
 DOCUMENT NUMBER: PREV200300546092  
 TITLE: Genetic analysis of high frequency rugose exopolysaccharide  
 production (HFRP) in epidemic Vibrio cholerae.  
 AUTHOR(S): Rashid, M. H. [Reprint Author]; Ali, A. [Reprint Author];  
**Karaolis, D. K. R.** [Reprint Author]

CORPORATE SOURCE: University of Maryland School of Medicine, Baltimore, MD, USA

SOURCE: Abstracts of the General Meeting of the American Society for Microbiology, (2003) Vol. 103, pp. J-015.  
<http://www.asmtusa.org/mtgsrc/generalmeeting.htm>. cd-rom.  
 Meeting Info.: 103rd American Society for Microbiology General Meeting. Washington, DC, USA. May 18-22, 2003.  
 American Society for Microbiology.  
 ISSN: 1060-2011 (ISSN print).

DOCUMENT TYPE: Conference; (Meeting)  
 Conference; Abstract; (Meeting Abstract)

LANGUAGE: English

ENTRY DATE: Entered STN: 19 Nov 2003  
 Last Updated on STN: 19 Nov 2003

AB *Vibrio cholerae* is the causative organism of the severe life threatening epidemic diarrheal disease called cholera. Some strains of *V. cholerae* can shift to a rugose colony phenotype whereby the cells produce copious amounts of exopolysaccharide (EPS). The production of this EPS promotes the persistence of *V. cholerae* as it promotes **biofilm** formation and increased resistance to various stresses such as acid, chlorine, UV light, and complement. Recently, we have shown that certain strains of *V. cholerae* can shift at high frequency to produce EPS in a process we called high frequency rugose EPS production (HFRP). We also found that HFRP was more common in epidemic *V. cholerae* strains than in non-pathogenic strains suggesting that EPS production and HFRP is important in epidemic strains and might provide these strains with an adaptive advantage in certain niches. In order to explore the genetic basis of HFRP in epidemic strains, we generated 10,000 mini-Tn5 mutants in the smooth 7th pandemic (El Tor) strain N16961 using minitransposon, pUT Km-2. The mutants were screened in order to identify mutants that were defective in their ability to switch to HFRP. A total of 29 mutants were found to be defective for HFRP. Sequencing and further analysis indicates that EPS biosynthetic genes and several regulatory genes have important roles in the genetic switch to HFRP. Our results suggest that HFRP is linked to a complex regulatory circuit in *V. cholerae*.

L58 ANSWER 24 OF 25 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on STN

ACCESSION NUMBER: 2003:517549 BIOSIS Full-text

DOCUMENT NUMBER: PREV200300520027

TITLE: The *Vibrio* pathogenicity island-encoded Mop protein modulates cholera toxin production and **biofilm** formation in epidemic *V. cholerae*.

AUTHOR(S): Zhang, D. [Reprint Author]; Sun, W. [Reprint Author]; **Karaolis, D.** [Reprint Author]

CORPORATE SOURCE: University of Maryland School of Medicine, Baltimore, MD, USA

SOURCE: Abstracts of the General Meeting of the American Society for Microbiology, (2003) Vol. 103, pp. B-301.  
<http://www.asmtusa.org/mtgsrc/generalmeeting.htm>. cd-rom.  
 Meeting Info.: 103rd American Society for Microbiology General Meeting. Washington, DC, USA. May 18-22, 2003.  
 American Society for Microbiology.  
 ISSN: 1060-2011 (ISSN print).

DOCUMENT TYPE: Conference; (Meeting)  
 Conference; Abstract; (Meeting Abstract)

LANGUAGE: English

ENTRY DATE: Entered STN: 5 Nov 2003  
 Last Updated on STN: 5 Nov 2003



AB Epidemic cholera is a severe diarrheal disease caused by specific toxigenic strains of *Vibrio cholerae*. These strains possess an essential virulence cluster called the *Vibrio* pathogenicity island (VPI) that is typically absent from nonpathogenic strains. The VPI is 41.2-Kb in size and encodes 29 potential proteins, several of which have no known function. In order to better understand the pathogenesis of epidemic *V. cholerae* strains, we are characterizing the role of the VPI and its genes in virulence. We report that the VPI-encoded Mop protein is a predicted 34-kDa periplasmic protein which has no homolog in the database and contains a zinc metalloprotease motif. We constructed a mop mutation in *V. cholerae* 7th pandemic (El Tor) strain 3083 and found that Mop has no role in the expression of TcpA and hemagglutinin protease (HAP) or in motility, however, a Mop mutant is significantly attenuated in cholera toxin expression and increased in **biofilm** formation. Mop appears to be a protease that modulates the expression of several secreted proteins in *V. cholerae*. Our studies also suggest that there are differences between *V. cholerae* strains in the phenotypes that are modulated by Mop. Our in vitro studies support our previous in vivo results which showed that Mop is involved in the virulence of epidemic *V. cholerae* strains.

L58 ANSWER 25 OF 25 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on STN

ACCESSION NUMBER: 2002:597257 BIOSIS Full-text

DOCUMENT NUMBER: PREV200200597257

TITLE: Analysis of the genetic switch for phenotypic conversion between the smooth and rugose exopolysaccharide phenotypes of *Vibrio cholerae*.

AUTHOR(S): Rashid, M. H. [Reprint author]; Ali, A. [Reprint author]; **Karaolis, D. K. R.** [Reprint author]

CORPORATE SOURCE: University of Maryland School of Medicine, Baltimore, MD, USA

SOURCE: Abstracts of the General Meeting of the American Society for Microbiology, (2002) Vol. 102, pp. 263. print. Meeting Info.: 102nd General Meeting of the American Society for Microbiology. Salt Lake City, UT, USA. May 19-23, 2002. American Society for Microbiology. ISSN: 1060-2011.

DOCUMENT TYPE: Conference; (Meeting)  
Conference; Abstract; (Meeting Abstract)

LANGUAGE: English

ENTRY DATE: Entered STN: 20 Nov 2002

Last Updated on STN: 20 Nov 2002

AB Cholera is a life threatening diarrheal disease caused by the bacterium *Vibrio cholerae*. Although the ability of epidemic *V. cholerae* strains to persist in the environment is critical in the epidemiology of cholera, the mechanisms underlying the endemicity and long-term persistence of epidemic *V. cholerae* strains are not well understood. Exopolysaccharide (EPS) production is important in **biofilm** formation and the persistence of many bacterial species. *V. cholerae* strains can under some conditions express a "rugose" colony phenotype due to copious production of EPS and which is thought to be important in the long-term persistence of the organism. We have recently discovered a phenomenon called "high frequency rugose EPS production" (HFRP) is unique to epidemic *V. cholerae* strains in which a high frequency of smooth cells convert to rugose cells expressing copious EPS. We also found HFRP strains with a high rate of switching from the rugose to smooth phenotype. These findings suggest that EPS and HFRP are important in epidemic *V. cholerae* strains, however, the genetics and physiology of rugose EPS production are not well understood. Transposon mutagenesis and further genetic analysis was used to identify the genes involved in the genetic switch controlling the phenotypic shift between smooth and rugose phenotypes. Several mutants that

are defective in this phenotypic conversion have been identified. Since EPS is important for long-term persistence and since HFRP is unique to epidemic V. cholerae strains, our findings provide valuable information regarding the genetic switch of a phenomenon that appears to be important in the epidemiology of epidemic V. cholerae strains.

**Text search history**

=&gt; d his L31

(FILE 'HCAPLUS' ENTERED AT 16:14:27 ON 19 OCT 2009)

L31 16 S L26 OR L30

=&gt; d que L31

L3 1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON CYCLIC GMP/CN  
 L4 24 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON ?CYCL?/CNS (L)  
 ?GUANOS?/CNS (L) ?MONOPHOSPHAT?/CNS  
 L5 457 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON ?CYCL?/CNS (L)  
 ?DIGUAN?/CNS  
 L6 1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON L5 AND ?ACID?/CNS  
 L7 2 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON "C-DI-GMP"/ONS  
 L12 7 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON "CYCLIC DIGUANYLATE"/  
 ONS  
 L16 7 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON "GUANOSINE 3',5'-CYCL  
 IC MONOPHOSPHATE"/ONS  
 L17 1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON "3'-5'-CYCLIC  
 DIGUANYLIC ACID"/ONS  
 L19 1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON "CYCLIC DIGUANYLIC  
 ACID"/ONS  
 L21 40435 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON (L3 OR L4 OR L5 OR L6  
 OR L7) OR L12 OR L16 OR L17 OR L19  
 L22 28335 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON ("CYCLIC GMP" OR  
 CYCLIC(W)GMP OR (CYCLIC(3W)DIGUANYL?) OR "C-GMP" OR "CGMP" OR  
 "C-DI-GMP" OR "BIS-(3',5') CYCLIC DIGUANYLIC ACID" OR "3',5'-CY  
 CLIC DI-GMP" OR "CYCLIC DINUCLEOTIDE DI-GMP" OR "BIS-(3'-5')-C  
 YCLIC DIMERIC GUANOSINE MONOPHOSPHATE" OR "C-(GPGP)" OR  
 "CGPGP")  
 L23 847 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON ( "GUANOSINE 3',5'-CYC  
 LIC MONOPHOSPHATE" OR "3'-5'-CYCLIC DIGUANYLIC ACID" OR  
 "BIS-(3'-5') CYCLIC DIGUANYLIC ACID" OR "CYCLIC DIGUANYLIC  
 ACID" OR "CYCLIC-BIS(3',5')DIGUANYLIC ACID")  
 L24 1837 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON (L21 OR L22 OR L23)  
 AND ((INHIBIT? OR REDUC? OR CONTROL? OR DIMINI? OR MINIM? OR  
 DELAY? OR RETARD? OR PREVENT? OR PROPHYL? OR ELIMIN? OR  
 DECREAS?) (5A) (MICROB? OR BACT? OR FUNG? OR PATHOGEN? OR  
 BIOFILM? OR BIO(W)FILM? OR BIOSLIM? OR BIO(W)SLIM? OR COLONI?  
 OR COLONY OR INFECT?))  
 L25 229 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L24 AND (SKIN? OR  
 DERM? OR EPIDERM? OR NASAL? OR NASO? OR PHARYN? OR SINUS? OR  
 SINO? OR MUCUS? OR MUCOS? OR MUCOUS? OR MEMBRAN?)  
 L26 10 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L25 AND ((MEDIC? OR  
 SURGIC? OR THERAP? OR PATIENT? OR TREAT? OR RECONSTRUCT? OR  
 ARTIFIC? OR REPLAC?) (5A) (DEVIC? OR IMPLEMENT? OR STENT? OR  
 CATHET? OR IMPLANT? OR PROSTHET? OR INDWELL? OR IN(W)DWELL?))  
 L27 71 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L24 AND (BIOFILM? OR  
 BIO(W)FILM? OR "IN VIVO" OR IN(W)VIVO? OR "IN VITRO" OR  
 IN(W)VITR?)  
 L28 55 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L27 AND ((BIOFILM? OR  
 BIO(W)FILM? OR BACT? OR PATHOGEN?) (3A) (INHIBIT? OR REDUC? OR  
 CONTROL? OR DIMINI? OR MINIM? OR MITIGAT? OR PREVENT? OR  
 STERIL? OR SANIT?))  
 L29 55 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L28 AND (BIOFILM? OR  
 BIO(W)FILM?)  
 L30 7 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L29 AND ((ADMINIST?  
 OR TREAT? OR APPLY? OR APPLICAT?) AND (PATIENT? OR MAMMAL? OR

10/565,591

SUBJECT? OR CLINIC?))

L31 16 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L26 OR L30

=> d his L40

(FILE 'HCAPLUS' ENTERED AT 16:41:51 ON 19 OCT 2009)

L40 3 S L39 AND ((ADMINIST? OR TREAT? OR APPLY? OR APPLICAT?) AND (PA

=> d que L40

L3 1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON CYCLIC GMP/CN

L4 24 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON ?CYCL?/CNS (L)  
?GUANOS?/CNS (L) ?MONOPHOSPHAT?/CNS

L5 457 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON ?CYCL?/CNS (L)  
?DIGUAN?/CNS

L6 1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON L5 AND ?ACID?/CNS

L7 2 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON "C-DI-GMP"/ONS

L12 7 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON "CYCLIC DIGUANYLATE"/  
ONS

L16 7 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON "GUANOSINE 3',5'-CYCL  
IC MONOPHOSPHATE"/ONS

L17 1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON "3'-5'-CYCLIC  
DIGUANYLIC ACID"/ONS

L19 1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON "CYCLIC DIGUANYLIC  
ACID"/ONS

L21 40435 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON (L3 OR L4 OR L5 OR L6  
OR L7) OR L12 OR L16 OR L17 OR L19

L34 47 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON (1044660-34-5/BI OR  
111-30-8/BI OR 132182-18-4/BI OR 132182-19-5/BI OR 132182-21-9/  
BI OR 132209-26-8/BI OR 132294-58-7/BI OR 146316-82-7/BI OR  
20816-12-0/BI OR 232933-52-7/BI OR 31348-80-8/BI OR 3353-33-1/B  
I OR 541-09-3/BI OR 548-62-9/BI OR 6018-53-7/BI OR 60307-63-3/B  
I OR 61093-23-0/BI OR 66-79-5/BI OR 75-56-9/BI OR 7681-52-9/BI  
OR 7722-84-1/BI OR 7782-50-5/BI OR 849214-01-3/BI OR 849214-02-  
4/BI OR 849214-03-5/BI OR 849214-04-6/BI OR 849214-05-7/BI OR  
849214-06-8/BI OR 849214-07-9/BI OR 849214-08-0/BI OR 849214-09  
-1/BI OR 849214-10-4/BI OR 849214-11-5/BI OR 849214-12-6/BI OR  
849214-13-7/BI OR 849214-14-8/BI OR 849214-15-9/BI OR 849214-16  
-0/BI OR 849447-99-0/BI OR 849448-00-6/BI OR 849448-01-7/BI OR  
849448-02-8/BI OR 849448-03-9/BI OR 9004-34-6/BI OR 9012-56-0/B  
I OR 9013-25-6/BI OR 9042-14-2/BI)

L35 2 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON L34 AND L21

L36 1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON 61093-23-0/RN

L37 3 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON 61093-23-0/CRN

L38 55 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON (L35 OR L36 OR L37)  
AND ((INHIBIT? OR REDUC? OR CONTROL? OR DIMINI? OR MINIM? OR  
DELAY? OR RETARD? OR PREVENT? OR PROPHYL? OR ELIMIN? OR  
DECREAS?) (5A) (MICROB? OR BACT? OR FUNG? OR PATHOGEN? OR  
BIOFILM? OR BIO(W)FILM? OR BIOSLIM? OR BIO(W)SLIM? OR COLONI?  
OR COLONY OR INFECT?))

L39 36 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L38 AND (BIOFILM? OR  
BIO(W)FILM? OR "BIOFILM")

L40 3 SEA FILE=HCAPLUS SPE=ON ABB=ON PLU=ON L39 AND ((ADMINIST?  
OR TREAT? OR APPLY? OR APPLICAT?) AND (PATIENT? OR MAMMAL? OR  
SUBJECT? OR CLINIC?))

=> d his L57

(FILE 'MEDLINE, BIOSIS, EMBASE, DRUGU' ENTERED AT 16:49:19 ON 19 OCT 2009)

```

L57          24 S L54 OR L56

=> d que L57
L3           1 SEA FILE=REGISTRY SPE=ON  ABB=ON  PLU=ON  CYCLIC GMP/CN
L5           457 SEA FILE=REGISTRY SPE=ON  ABB=ON  PLU=ON  ?CYCL?/CNS (L)
              ?DIGUAN?/CNS
L6           1 SEA FILE=REGISTRY SPE=ON  ABB=ON  PLU=ON  L5 AND ?ACID?/CNS
L7           2 SEA FILE=REGISTRY SPE=ON  ABB=ON  PLU=ON  "C-DI-GMP"/ONS
L12          7 SEA FILE=REGISTRY SPE=ON  ABB=ON  PLU=ON  "CYCLIC DIGUANYLATE"/
              ONS
L16          7 SEA FILE=REGISTRY SPE=ON  ABB=ON  PLU=ON  "GUANOSINE 3',5'-CYCL
              IC MONOPHOSPHATE"/ONS
L17          1 SEA FILE=REGISTRY SPE=ON  ABB=ON  PLU=ON  "3'-5'-CYCLIC
              DIGUANYLIC ACID"/ONS
L19          1 SEA FILE=REGISTRY SPE=ON  ABB=ON  PLU=ON  "CYCLIC DIGUANYLIC
              ACID"/ONS
L22          28335 SEA FILE=HCAPLUS SPE=ON  ABB=ON  PLU=ON  ("CYCLIC GMP" OR
              CYCLIC(W)GMP OR (CYCLIC(3W)DIGUANYL?) OR "C-GMP" OR "CGMP" OR
              "C-DI-GMP" OR "BIS-(3',5') CYCLIC DIGUANYLIC ACID" OR "3',5'-CY
              CLIC DI-GMP" OR "CYCLIC DINUCLEOTIDE DI-GMP" OR "BIS-(3'-5')-C
              YCLIC DIMERIC GUANOSINE MONOPHOSPHATE" OR "C-(GPGP)" OR
              "CGPGP")
L23          847 SEA FILE=HCAPLUS SPE=ON  ABB=ON  PLU=ON  ( "GUANOSINE 3',5'-CYC
              LIC MONOPHOSPHATE" OR "3'-5'-CYCLIC DIGUANYLIC ACID" OR
              "BIS-(3'-5') CYCLIC DIGUANYLIC ACID" OR "CYCLIC DIGUANYLIC
              ACID" OR "CYCLIC-BIS(3',5')DIGUANYLIC ACID")
L45          58517 SEA L3 OR L6 OR L7 OR L12 OR L16 OR L17 OR L19
L46          58359 SEA L45 AND (L22 OR L23)
L47          118 SEA L46 AND (BIOFILM? OR BIO(W) FILM? OR "BIOFILM")
L49          6 SEA L47 AND ((ADMINIST? OR TREAT? OR APPLY? OR APPLICAT?) AND
              (PATIENT? OR MAMMAL? OR SUBJECT? OR CLINIC?))
L50          44 SEA L47 AND ((INHIBIT? OR REDUC? OR CONTROL? OR DIMINI? OR
              MINIM? OR DELAY? OR RETARD? OR PREVENT? OR PROPHYL? OR ELIMIN?
              OR DECREAS?) (5N) (MICROB? OR BACT? OR FUNG? OR PATHOGEN? OR
              BIOFILM? OR BIO(W) FILM? OR BIOSLIM? OR BIO(W) SLIM? OR
              COLONI? OR COLONY OR INFECT?))
L51          15 SEA L47 AND (SKIN? OR DERM? OR EPIDERM? OR NASAL? OR NASO? OR
              PHARYN? OR SINUS? OR SINO? OR MUCUS? OR MUCOS? OR MUCOUS? OR
              MEMBRAN?)
L52          9 SEA L50 AND (L49 OR L51)
L53          10 SEA L50 AND (ADMINIST? OR PATIENT? OR MAMMAL? OR TREAT?)
L54          24 SEA L49 OR (L51 OR L52 OR L53)
L55          38 SEA L47 AND (STAPHYLOCOCC? OR VIBRIO? OR GONOCOCC?)
L56          10 SEA L54 AND L55
L57          24 SEA L54 OR L56

```

```

=> dup rem L31 L40 L57
PROCESSING COMPLETED FOR L31
PROCESSING COMPLETED FOR L40
PROCESSING COMPLETED FOR L57
L59          30 DUP REM L31 L40 L57 (13 DUPLICATES REMOVED)
              ANSWERS '1-16' FROM FILE HCAPLUS
              ANSWERS '17-28' FROM FILE MEDLINE
              ANSWERS '29-30' FROM FILE EMBASE

```

**Text search results: HCAPLUS**

=&gt; d L59 1-16 ibib ed abs hitrn

L59 ANSWER 1 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 5  
 ACCESSION NUMBER: 2005:300236 HCAPLUS Full-text  
 DOCUMENT NUMBER: 142:367640  
 TITLE: Method for attenuating virulence of **microbial pathogens** and **inhibiting microbial biofilm** formation by using **c-di-GMP** and cyclic dinucleotide analogs  
 INVENTOR(S): Karaolis, David K. R.  
 PATENT ASSIGNEE(S): University of Maryland, USA  
 SOURCE: PCT Int. Appl., 118 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005030186	A2	20050407	WO 2004-US23498	20040722
WO 2005030186	A3	20050714		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2004275696	A1	20050407	AU 2004-275696	20040722
CA 2533873	A1	20050407	CA 2004-2533873	20040722
EP 1651242	A2	20060503	EP 2004-809506	20040722
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK			
JP 2007500697	T	20070118	JP 2006-521912	20040722
US 20070244059	A1	20071018	US 2006- <b>565591</b>	20061006
PRIORITY APPLN. INFO.:			US 2003-490029P	P 20030728
			WO 2004-US23498	W 20040722

ED Entered STN: 07 Apr 2005

AB The present invention relates to the use of the cyclic dinucleotide **c-di-GMP** and cyclic dinucleotide analogs thereof in a method for attenuating virulence of a **microbial pathogen** or for **inhibiting** or **reducing colonization** by a **microbial pathogen**. This method further **inhibits microbial biofilm** formation and is capable of **treating** bacterial infections. The **microbial colonization** or **biofilm** formation **inhibited** or **reduced** may be on the **skin** or on **nasal** or **mucosal** surface. The **microbial colonization** or **biofilm** formation **inhibited** can also be on the surfaces of **medical devices**, especially those in close contact with the **patient**, as well on the surfaces of industrial and construction material where microbial colonization and **biofilm** formation is of concern.

IT **146316-82-7**

RL: PRPH (Prophetic)

10/565,591

(Method for attenuating virulence of **microbial pathogens** and **inhibiting microbial biofilm** formation by using **c-di-GMP** and cyclic dinucleotide analogs)

IT **61093-23-0D**, carboxy/phosphoalkylene ether derivs.

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(attenuating virulence of **microbial pathogens** and **inhibiting microbial biofilm** formation by using **c-di-GMP** and cyclic dinucleotide analogs)

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)

L59 ANSWER 2 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 6

ACCESSION NUMBER: 2005:887352 HCAPLUS Full-text

DOCUMENT NUMBER: 143:402354

TITLE: Aminoglycoside antibiotics induce bacterial **biofilm** formation

AUTHOR(S): Hoffman, Lucas R.; D'Argenio, David A.; MacCoss, Michael J.; Zhang, Zhaoying; Jones, Roger A.; Miller, Samuel I.

CORPORATE SOURCE: Department of Pediatrics, University of Washington, Seattle, WA, 98195, USA

SOURCE: Nature (London, United Kingdom) (2005), 436(7054), 1171-1175

CODEN: NATUAS; ISSN: 0028-0836

PUBLISHER: Nature Publishing Group

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 25 Aug 2005

AB **Biofilms** are adherent aggregates of bacterial cells that form on biotic and abiotic surfaces, including human tissues. **Biofilms** resist antibiotic **treatment** and contribute to bacterial persistence in chronic infections. Hence, the elucidation of the mechanisms by which **biofilms** are formed may assist in the **treatment** of chronic infections, such as *Pseudomonas aeruginosa* in the airways of **patients** with cystic fibrosis. Here we show that subinhibitory concns. of aminoglycoside antibiotics induce **biofilm** formation in *P. aeruginosa* and *Escherichia coli*. In *P. aeruginosa*, a gene, which we designated aminoglycoside response regulator (*arr*), was essential for this induction and contributed to **biofilm**-specific aminoglycoside resistance. The *arr* gene is predicted to encode an inner-membrane phosphodiesterase whose substrate is cyclic di-guanosine monophosphate (**c-di-GMP**)-a bacterial second messenger that regulates cell surface adhesiveness. We found that membranes from *arr* mutants had diminished **c-di-GMP** phosphodiesterase activity, and *P. aeruginosa* cells with a mutation changing a predicted catalytic residue of *Arr* were defective in their **biofilm** response to tobramycin. Furthermore, tobramycin-inducible **biofilm** formation was **inhibited** by exogenous GTP, which is known to inhibit **c-di-GMP** phosphodiesterase activity. Our results demonstrate that **biofilm** formation can be a specific, defensive reaction to the presence of antibiotics, and indicate that the mol. basis of this response includes alterations in the level of **c-di-GMP**.

IT **57-92-1**, Streptomycin, biological studies **61093-23-0**  
**338732-46-0**

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(aminoglycoside antibiotics induce bacterial **biofilm** formation)

OS.CITING REF COUNT: 95 THERE ARE 95 CAPLUS RECORDS THAT CITE THIS  
RECORD (95 CITINGS)

REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L59 ANSWER 3 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 7

ACCESSION NUMBER: 2005:229578 HCAPLUS Full-text

DOCUMENT NUMBER: 142:426617

TITLE: **c-di-GMP (3'-5'-cyclic diguanylic acid)** inhibits *Staphylococcus aureus* cell-cell interactions and **biofilm** formation

AUTHOR(S): Karaolis, David K. R.; Rashid, Mohammed H.; Chythanya, Rajanna; Luo, Wensheng; Hyodo, Mamoru; Hayakawa, Yoshihiro

CORPORATE SOURCE: Department of Epidemiology and Preventive Medicine, University of Maryland School of Medicine, Baltimore, MD, 21201, USA

SOURCE: Antimicrobial Agents and Chemotherapy (2005), 49(3), 1029-1038

CODEN: AMACCQ; ISSN: 0066-4804

PUBLISHER: American Society for Microbiology

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 17 Mar 2005

AB *Staphylococcus aureus* is an important pathogen of humans and animals, and antibiotic resistance is a public health concern. **Biofilm** formation is essential in virulence and pathogenesis, and the ability to resist antibiotic **treatment** results in difficult-to- **treat** and persistent infections. As such, novel antimicrobial approaches are of great interest to the scientific, medical, and agriculture communities. We recently proposed that modulating levels of the cyclic dinucleotide signaling mol., **c-di- GMP (cyclic diguanylate [3', 5'-cyclic diguanylic acid], cGpGp)**, has utility in regulating phenotypes of prokaryotes. We report that extracellular **c-di-GMP** shows activity against human **clin.** and bovine intramammary mastitis isolates of *S. aureus*, including methicillin-resistant *S. aureus* (MRSA) isolates. We show that chemical synthesized **c-di- GMP** is soluble and stable in water and physiol. saline and stable following boiling and exposure to acid and alkali. **Treatment** of *S. aureus* with extracellular **c-di-GMP** inhibited cell-to-cell (intercellular) adhesive interactions in liquid medium and **reduced** (>50%) **biofilm** formation in human and bovine isolates compared to untreated controls. **C- di-GMP** inhibited the adherence of *S. aureus* to human epithelial HeLa cells. The cyclic nucleotide analogs **cGMP** and **cAMP** had a lesser **inhibitory** effect on **biofilms**, while 5'-GMP had no major effect. We propose that cyclic dinucleotides such as **c-di-GMP**, used either alone or in combination with other antimicrobial agents, represent a novel and attractive approach in the development of intervention strategies for the **prevention** of **biofilms** and the **control** and **treatment** of **infection**.

IT 61093-23-0

RL: BSU (Biological study, unclassified); BIOL (Biological study)

(**c-di-GMP (3'-5'-cyclic diguanylic acid)** inhibits *Staphylococcus aureus* cell-cell interactions and **biofilm** formation)

OS.CITING REF COUNT: 30 THERE ARE 30 CAPLUS RECORDS THAT CITE THIS RECORD (30 CITINGS)

REFERENCE COUNT: 64 THERE ARE 64 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L59 ANSWER 4 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2009:1139004 HCAPLUS Full-text

DOCUMENT NUMBER: 151:373900

TITLE: Use of ellagitannins as **inhibitors** of **bacterial** quorum sensing



10/565,591

INVENTOR(S): Mathee, Kalai; Adonizio, Allison L.; Ausubel, Frederick; Clardy, Jon; Bennett, Bradley; Downum, Kelsey  
 PATENT ASSIGNEE(S): The Florida International Uinversity, USA  
 SOURCE: PCT Int. Appl., 59pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
WO 2009114810	A2	20090917	WO 2009-US37163	20090313
W: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				

PRIORITY APPLN. INFO.: US 2008-36812P P 20080314

ED Entered STN: 18 Sep 2009

AB The invention provides materials and methods for the **inhibition** of **bacterial** quorum sensing (QS). The invention also provides methods for treating bacterial infections by administration of one or more ellagitannins in amount effective to **inhibit bacterial QS**.

IT **57-92-1**, Streptomycin **57-92-1D**, Streptomycin, derivs.

RL: PAC (Pharmacological activity); TEM (Technical or engineered material use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (ellagitannins as **inhibitors** of **bacterial** quorum sensing, use in treatment of bacterial infection, and use with other agents)

L59 ANSWER 5 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2009:1080981 HCAPLUS Full-text

DOCUMENT NUMBER: 151:329919

TITLE: Recombinant **bacteriophage** containing nucleotides for **inhibiting** antibiotic resistance, survival, SOS response and/or defense genes, and their use in combination of antimicrobial agents in **eliminating bacteria** in animals or surfaces

INVENTOR(S): Collins, James J.; Lu, Timothy Kuan-Ta

PATENT ASSIGNEE(S): Boston University, USA; Massachusetts Institute of Technology

SOURCE: PCT Int. Appl., 229pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

WO 2009108406	A2	20090903	WO 2009-US30755	20090112
W: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				

PRIORITY APPLN. INFO.:

US 2008-20197P

P 20080110

ED Entered STN: 04 Sep 2009

AB The invention provides recombinant, genetically engineered bacteriophages, and their use in combination with an antimicrobial agent to **reduce** or **eliminate bacteria** in animals (including humans), and/or on surfaces. Specifically, recombinant bacteriophages, developed from M13 or T7, contain nucleic acids designed to: (a) **inhibit** at least one **bacterial** antibiotic resistance gene (such as cat, vanA or mecD) and/or survival gene (such as RecA, RecB, RecC, spot or RelA); (b) repress at least one SOS response or defense gene; and/or (c) express a protein (such as porin) that increases susceptibility of a bacterial cell to an antimicrobial agent, wherein said protein modifies a specific pathway. The invention also provides compns. and kits comprising said recombinant bacteriophages and at least on antimicrobial agent.

IT **57-92-1D**, Streptomycin, variant and/or analog thereof

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(bacteriophage containing nucleotides for **inhibiting**antibiotic resistance, survival, SOS response and/or defense genes, and their use in combination of antimicrobial agents in **eliminating bacteria** in animals or surfaces)

L59 ANSWER 6 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2008:416834 HCAPLUS Full-text

DOCUMENT NUMBER: 148:397490

TITLE: Analyte sensing device with external control unit and implantable biosensor for continuously monitoring metabolic levels of analytes

INVENTOR(S): Grantham, Daniel H.; Jain, Faquir;

Papadimitrakopoulos, Fotios; Burgess, Diane

PATENT ASSIGNEE(S): University of Connecticut, USA; Optoelectronics

Systems Consulting, Inc. "Osci"; Grantham, G.,

Deborah; Salisbury, Jane

SOURCE: PCT Int. Appl., 76pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2008039543	A1	20080403	WO 2007-US21042	20070927
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CH, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL,				

10/565,591

PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN,  
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW  
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,  
IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF,  
BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW,  
GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ,  
BY, KG, KZ, MD, RU, TJ, TM

US 20080154101 A1 20080626 US 2007-862866 20070927  
EP 2079358 A1 20090722 EP 2007-839072 20070927

R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,  
IS, IT, LI, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR

PRIORITY APPLN. INFO.: US 2006-827104P P 20060927  
WO 2007-US21042 W 20070927

ED Entered STN: 03 Apr 2008

AB Disclosed herein is an analyte sensing device capable of continuously monitoring metabolic levels of a plurality of analytes. The device comprises an external unit, which, for example, could be worn around the wrist like a wristwatch or could be incorporated into a cell phone or PDA device, and an implantable sensor platform that is suitable, for example, for implantation under the **skin**. The external device and the internal device are in wireless communication. In one embodiment, the external device and the internal device are operationally linked by a feedback system. In one embodiment, the internal device is encapsulated in a biocompatible coating capable of controlling the local tissue environment in order to prevent/minimize inflammation and fibrosis, promote neo-angiogenesis and wound healing and this facilitate device functionality.

IT **57-92-1**, Streptomycin

RL: TEM (Technical or engineered material use); THU (Therapeutic use);  
BIOL (Biological study); USES (Uses)

(analyte sensing device with external control unit and implantable biosensor for continuously monitoring metabolic levels of analytes)

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L59 ANSWER 7 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2008:160537 HCAPLUS Full-text

DOCUMENT NUMBER: 148:209646

TITLE: Plate for selection of antibiotics against  
**biofilm** infections

INVENTOR(S): Olson, Merle E.; Ceri, Howard

PATENT ASSIGNEE(S): MBEC Bioproducts Inc., Can.

SOURCE: PCT Int. Appl., 59pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	---	-----	-----	-----
WO 2008014581	A1	20080207	WO 2006-CA1226	20060724
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ,			

10/565,591

CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,  
KG, KZ, MD, RU, TJ, TM

CA 2616526 A1 20070122 CA 2006-2616526 20060724  
US 20080318269 A1 20081225 US 2008-996480 20080819  
PRIORITY APPLN. INFO.: US 2005-701858P P 20050722  
WO 2006-CA1226 W 20060724

ED Entered STN: 08 Feb 2008

AB This invention is a diagnostic plate that can be used to select antibiotic combinations with efficacy against microorganisms growing as a **biofilm**. The plate allows growth of **biofilm** on a plurality of projections, and the subsequent simultaneous challenge of **biofilms** on all projections of the plate to independent concns. and combinations of anti-**biofilm** agents. Resistance of microorganisms to antibiotics is higher when they grow as a **biofilm**, as compared to when they grow in a planktonic state which is usually used to determine their level of antibiotic sensitivity. Growth of microorganisms that slough off the **biofilm** in the anti- **biofilm** agent challenge dets. the Min. **Inhibitory** Concentration (MIC) which relates to sensitivity of the microorganisms in a planktonic state. Growth of any surviving microorganisms from the **biofilm** in a subsequent recovery step dets. the **Minimal Biofilm** Eradication Concentration (MBEC) which relates to the sensitivity of the microorganisms growing as a **biofilm**. Enumeration of the surviving microorganisms in the recovery step dets. the Min. Biocidal Concentration (MBC). A Staphylococcus test plate was developed to evaluate antibiotics and antibiotic combinations against Staphylococcus aureus **biofilms**. The 96-well plate had gentamicin, clindamycin, cefazolin, cloxacillin, rifampin, vancomycin, linezolid, ampicillin sublactam, ciprofloxacin, various combinations of the antibiotics, a growth control, and a sterility control. The sensitivity of planktonic and **biofilm** forms of S. aureus to individual and combination agents could be determined within about 48 h. S. aureus was sensitive to multiple antibiotics and antibiotic combinations as planktonic forms, but was significantly more resistant as a **biofilm**.

IT 57-92-1

RL: BSU (Biological study, unclassified); PAC (Pharmacological activity);  
PRPH (Prophetic); THU (Therapeutic use); BIOL (Biological study); USES  
(Uses)

(plate for selection of antibiotics and fungicides against  
**biofilm** infections)

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L59 ANSWER 8 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2008:157144 HCAPLUS Full-text

DOCUMENT NUMBER: 148:209645

TITLE: Plate for selection of antibiotics against  
**biofilm** infections

INVENTOR(S): Olson, Merle E.; Ceri, Howard

PATENT ASSIGNEE(S): MBEC Bioproducts Inc., Can.

SOURCE: PCT Int. Appl., 64pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
WO 2008014580	A1	20080207	WO 2006-CA1218	20060724
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,			

GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

CA 2616559 A1 20080124 CA 2006-2616559 20060724  
 EP 1915458 A1 20080430 EP 2006-761179 20060724

R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, RS

KR 2008081891 A 20080910 KR 2008-704354 20080222  
 CN 101283104 A 20081008 CN 2006-80030645 20080222  
 US 20080318268 A1 20081225 US 2008-996478 20080819

PRIORITY APPLN. INFO.: US 2005-701858P P 20050722  
 WO 2006-CA1218 W 20060724

ED Entered STN: 07 Feb 2008

AB This invention is a diagnostic plate that can be used to select antibiotic combinations with efficacy against microorganisms growing as a **biofilm**. The plate allows growth of **biofilm** on a plurality of projections, and the subsequent simultaneous challenge of **biofilms** on all projections of the plate to independent concns. and combinations of anti-**biofilm** agents. Resistance of microorganisms to antibiotics is higher when they grow as a **biofilm**, as compared to when they grow in a planktonic state which is usually used to determine their level of antibiotic sensitivity. Growth of microorganisms that slough off the **biofilm** in the anti- **biofilm** agent challenge dets. the Min. **Inhibitory** Concentration (MIC) which relates to sensitivity of the microorganisms in a planktonic state. Growth of any surviving microorganisms from the **biofilm** in a subsequent recovery step dets. the **Minimal Biofilm** Eradication Concentration (MBEC) which relates to the sensitivity of the microorganisms growing as a **biofilm**. Enumeration of the surviving microorganisms in the recovery step dets. the Min. Biocidal Concentration (MBC). A Staphylococcus test plate was developed to evaluate antibiotics and antibiotic combinations against Staphylococcus aureus **biofilms**. The 96-well plate had gentamicin, clindamycin, cefazolin, cloxacillin, rifampin, vancomycin, linezolid, ampicillin sublactam, ciprofloxacin, various combinations of the antibiotics, a growth control, and a sterility control. The sensitivity of planktonic and **biofilm** forms of S. aureus to individual and combination agents could be determined within about 48 h. S. aureus was sensitive to multiple antibiotics and antibiotic combinations as planktonic forms, but was significantly more resistant as a **biofilm**.

IT **57-92-1**, Streptomycin  
 RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); PRPH (Prophetic); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (plate for selection of antibiotics and fungicides against **biofilm** infections)

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
 REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L59 ANSWER 9 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2007:1278450 HCAPLUS Full-text  
 DOCUMENT NUMBER: 147:508656  
 TITLE: Antimicrobial site dressings comprising, for example,

10/565,591

a silver compound for use with percutaneous  
**medical devices**

INVENTOR(S): McMaken, Jack D.; Gibbins, Bruce L.  
PATENT ASSIGNEE(S): Acrymed, Inc., USA  
SOURCE: PCT Int. Appl., 22pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
WO 2007127236	A2	20071108	WO 2007-US9997	20070425
WO 2007127236	A3	20081106		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA				
US 20070293800	A1	20071220	US 2007-789701	20070425
EP 2015722	A2	20090121	EP 2007-755996	20070425
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, RS				
PRIORITY APPLN. INFO.:			US 2006-796039P	P 20060428
			WO 2007-US9997	W 20070425

ED Entered STN: 09 Nov 2007

AB The present invention comprises antimicrobial articles for use with a percutaneous device, comprising a matrix which may contact the percutaneous device in a three-dimensional mode and release antimicrobial agents (e.g., silver ions) to the percutaneous device access site. In addition, the antimicrobial article of the present invention may donate moisture to a dry **dermal** site (e.g., a dry wound bed) and/or absorb liquid or exudates of a **dermal** site. The present invention also comprises methods for treating and/or **preventing an infection** using the antimicrobial articles of the present invention. Thus, a silver-containing polyacrylate matrix was made by mixing 185 g acrylamide and 2 g bisacrylamide into 3330 g of water containing 33.3 g of sodium chloride. To this mixture, was added 21 g of guar gum and 188 g of glycerol. After mixing to homogeneity, a solution containing 0.563 g silver nitrate was slowly added to the mixing batch. The polymerization of the mixture was accomplished by blending 1.8 mL TEMED and 2.6 g ammonium persulfate into the mixture. The mixture was poured into the appropriate molds before polymerization in a dark place. The gelled polymer was removed from the mold, dehydrated by mild heat and then rehydrated by humidification to a desired moisture content, 22% weight/weight. The matrix was then cut to form the article with one or two passages for use with percutaneous devices.

IT **57-92-1**, Streptomycin

RL: TEM (Technical or engineered material use); THU (Therapeutic use);  
BIOL (Biological study); USES (Uses)  
(antimicrobial site dressings for **prevention of infection** related to use of percutaneous **medical devices**)

10/565,591

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)

L59 ANSWER 10 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN  
 ACCESSION NUMBER: 2007:675678 HCAPLUS Full-text  
 DOCUMENT NUMBER: 147:87617  
 TITLE: Use of rifamycins for **treatment** of  
 infections, including **prosthetic** joint  
 infections  
 INVENTOR(S): Murphy, Christopher K.; Rothstein, David M.  
 PATENT ASSIGNEE(S): Activbiotics, Inc., USA  
 SOURCE: PCT Int. Appl., 98pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	---	-----	-----	-----
WO 2007070084	A1	20070621	WO 2006-US18559	20060515
W:				
AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,				
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,				
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR,				
KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX,				
MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE,				
SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC,				
VN, YU, ZA, ZM, ZW				
RW:				
AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,				
IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ,				
CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH,				
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,				
KG, KZ, MD, RU, TJ, TM				
AU 2006325493	A1	20070621	AU 2006-325493	20060515
CA 2631954	A1	20070621	CA 2006-2631954	20060515
EP 1971342	A1	20080924	EP 2006-759753	20060515
R:				
AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,				
IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR				
US 20070142392	A1	20070621	US 2006-638738	20061214
IN 2008DN05060	A	20080926	IN 2008-DN5060	20080612
MX 2008007809	A	20080915	MX 2008-7809	20080613
CN 101365455	A	20090211	CN 2006-80052335	20080805
PRIORITY APPLN. INFO.:			US 2005-750774P	P 20051215
			WO 2006-US18559	W 20060515

ED Entered STN: 22 Jun 2007

AB The invention provides methods, compns., and kits using rifamycin compds.  
 (Markush included) for treating a variety of bacterial infections, including  
**prosthetic** joint infections, infections caused by **medical implants**, infectious  
 arthritis, and osteomyelitis. The rifamycin compds. may be used in  
 conjunction with other antibiotics.

IT **57-92-1**, Streptomycin

RL: PAC (Pharmacological activity); TEM (Technical or engineered material  
 use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (rifamycins for **treatment** of infections, including  
**prosthetic** joint infections, and use with other antibiotics)

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L59 ANSWER 11 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2008:169442 HCAPLUS Full-text

DOCUMENT NUMBER: 148:513140

TITLE: Effect of different antimicrobial agents on Staphylococcus aureus adhesiveness and **biofilm** formation

AUTHOR(S): Yassien, M.; Al-Ansari, S.

CORPORATE SOURCE: Faculty of Pharmacy, King Abdul Aziz University, Jeddah, Saudi Arabia

SOURCE: New Egyptian Journal of Microbiology (2006), 13, 29-51  
CODEN: NEJMCI; ISSN: 1687-1219

PUBLISHER: Egyptian Society for Biotechnology

DOCUMENT TYPE: Journal

LANGUAGE: English

ED Entered STN: 11 Feb 2008

AB The effect of fluoroquinolones (ciprofloxacin, ofloxacin, and levofloxacin), clindamycin,  $\beta$ -lactams (cefoperazone, cefotaxime, cefepime), streptomycin, and vancomycin on the adherence and **biofilm** formation by Staphylococcus aureus (12 **clin.** isolates) was studied. In the presence of 1/2 MIC, 1/4 MIC and 1/8 MIC, the optical d. of the formed **biofilms** on plastic surfaces was **reduced** to 24-59.9%, 32.4-76.7% and 49.7-88.5% of the controls, resp. **Treatment** of the preformed **biofilms** with high concns. (25-200  $\mu\text{g/mL}$ ) of the tested agents caused reduction in the optical d. of the adherent **biofilms** to a range from 52.3 to 87.7% of the control. In an in-vitro model of vascular catheter **colonization**, the tested subinhibitory concns. **reduced** the percentage of the viable adherent cells to 32.1-71.6%, 42.5-85.6%, and 60.3-95.3% of the controls, resp. The tested fluoroquinolones and clindamycin are significantly more active than the other tested agents, and levofloxacin was the most active one. The vascular catheter segments precolonized with S. aureus for 24 h and exposed to 50  $\mu\text{g/mL}$  (4-31 times MIC) of the tested fluoroquinolones and clindamycin for 2 h showed few viable adherent cells (7-13 CFU/segment), while no adherent viable cells were cultured in the presence of 100  $\mu\text{g/mL}$  (8-62 times MIC). Also, the tested subinhibitory concns. **reduced** the percentage of the viable **bacterial** cells adherent to the surface of human lung epithelial A549 cells to the range of 30.1-79.2%, 41.1-89.3%, and 60.9-96.2% of the control, resp. **Treatment** of the A549 cells, preattached with bacterial cells, with the tested agents at concns. of 5, 10, and 20  $\mu\text{g/mL}$  (1/4-50 times MIC) reduced the range of the percentage of the adherent cells to 53.2-88.3%, 33.8-79.2%, and 27.2-68.1% of the control, resp. The superior activity of the tested fluoroquinolones and clindamycin was also observed. The obtained data show that subinhibitory concns. of ciprofloxacin, ofloxacin, levofloxacin, and clindamycin efficiently **reduced** the **biofilm** formation and adherence of S. aureus to the surfaces of plastics, vascular catheters, and human lung epithelial A549 cells. Also, higher concns. ( $\geq$  MIC) of fluoroquinolones and clindamycin were able to eradicate the adherent S. aureus.

IT 57-92-1, Streptomycin

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(effect of different antimicrobial agents on Staphylococcus aureus adhesiveness and **biofilm** formation)

REFERENCE COUNT: 63 THERE ARE 63 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L59 ANSWER 12 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2005:1126608 HCAPLUS Full-text

DOCUMENT NUMBER: 143:393137

TITLE: Novel modification of medical prostheses by coating with therapeutic agents

INVENTOR(S): Mansouri, Mohammad David; Darouiche, Rabih O.



PATENT ASSIGNEE(S): Baylor College of Medicine, USA  
 SOURCE: PCT Int. Appl., 37 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005096990	A2	20051020	WO 2005-US10944	20050331
WO 2005096990	A3	20070208		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2005231417	A1	20051020	AU 2005-231417	20050331
CA 2561496	A1	20051020	CA 2005-2561496	20050331
US 20050271694	A1	20051208	US 2005-95975	20050331
US 7238363	B2	20070703		
EP 1737378	A2	20070103	EP 2005-731606	20050331
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, LV, MK, YU			
PRIORITY APPLN. INFO.:			US 2004-558918P	P 20040402
			WO 2005-US10944	W 20050331

ED Entered STN: 20 Oct 2005

AB The incorporation of one or more therapeutic agents on metallic and non-metallic medical prostheses is provided. The therapeutic agent can be used, for example, to **prevent**, treat, or **reduce bacterial and fungal infections** associated with these implants. Addnl., the therapeutic agents can be used to effect other therapeutic benefits. Specifically, a bilayer therapeutic coating is applied in two steps. Addnl., non-antimicrobial therapeutic agents may be incorporated in this coating to treat, prevent, modify, or stimulate certain clin. bioactivities. Titanium cylinders were coated with a solution containing non-chromated hide powder, rifampin, and minocycline in glacial acetic acid. Antimicrobial activity of the coated device against *Staphylococcus epidermidis*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Escherichia coli*, and *Candida albicans* was shown.

IT **57-92-1**, Streptomycin, biological studies

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (novel modification of medical prostheses by coating with therapeutic agents)

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L59 ANSWER 13 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2004:412787 HCAPLUS Full-text

DOCUMENT NUMBER: 140:395549

TITLE: Controlled and continued delivery of rifaximin and/or

other substances  
 INVENTOR(S): Chiarelli, Piero; Dalseno, Renzo  
 PATENT ASSIGNEE(S): Italy  
 SOURCE: PCT Int. Appl., 15 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004041240	A1	20040521	WO 2003-EP12346	20031105
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003276254	A1	20040607	AU 2003-276254	20031105
EP 1560566	A1	20050810	EP 2003-810439	20031105
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
US 20060110447	A1	20060525	US 2005-533768	20050504
PRIORITY APPLN. INFO.:			IT 2002-FI212	A 20021105
			IT 2002-MI2438	A 20021118
			IT 2003-PI5	A 20030114
			IT 2003-PI13	A 20030221
			WO 2003-EP12346	W 20031105

ED Entered STN: 21 May 2004

AB A gum-like device is designed for the controlled and continued delivery of rifaximin, without producing the usually intense red coloration, for the resolution of the **infections** and the **reduction** of the inflammation in the oral cavity and in the laryngo-pharyngeal one. The device also protects either the gum or the dental apparatus from acute infections, from the infiltration and the stagnation of the food, and fights chronic infections such as in the periodontal pockets. Moreover, the device can be used to protect the gum from the traumatizing collision that the food exercises during the mastication.

IT **57-92-1**, Streptomycin, biological studies

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (gum-like device for controlled and continued delivery of rifaximin and other substances)

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L59 ANSWER 14 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2002:977651 HCAPLUS Full-text

DOCUMENT NUMBER: 138:61381

TITLE: Biofilm degradation or sloughing compositions containing furanones

INVENTOR(S): Kjelleberg, Staffan; Givskov, Michael; Hentzer, Morten

PATENT ASSIGNEE(S): Unisearch Limited, Australia

SOURCE: PCT Int. Appl., 69 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002102370	A1	20021227	WO 2002-AU797	20020618
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2002312648	A1	20030102	AU 2002-312648	20020618
US 20040147595	A1	20040729	US 2004-481250	20040331
PRIORITY APPLN. INFO.:			AU 2001-5754	A 20010618
			WO 2002-AU797	W 20020618

OTHER SOURCE(S): MARPAT 138:61381

ED Entered STN: 29 Dec 2002

AB The present invention relates to a method for the regulation and **control** of **biofilm** layers. In particular, the present invention is concerned with methods for degrading or causing sloughing of biofilms from surfaces (e.g., **medical** goods, **implants**, household furnishings, cooling systems in power plants). The invention is also related to compns. suitable for use in carrying out these methods. Thus, halogenated furanones were tested 8 different concns. The inhibitory activity of each compound on the fluorescent phenotype was diminished as the concentration increased.

IT **57-92-1**, Streptomycin, biological studies

RL: PAC (Pharmacological activity); BIOL (Biological study)

(biofilm degradation or sloughing compns. containing furanones)

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L59 ANSWER 15 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2002:241243 HCAPLUS Full-text

DOCUMENT NUMBER: 136:284492

TITLE: Compositions for treating biofilm

INVENTOR(S): Budny, John A.; Budny, Matthew J.

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 12 pp., Cont.-in-part of U. S. Ser. No. 587,818.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 7

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20020037260	A1	20020328	US 2001-876248	20010606
US 5871714	A	19990216	US 1997-951393	19971016
US 6159447	A	20001212	US 1999-249674	19990212
US 6830745	B1	20041214	US 2000-587818	20000606
PRIORITY APPLN. INFO.:			US 1997-951393	A2 19971016

US 1999-249674

A2 19990212

US 2000-587818

A2 20000606

ED Entered STN: 28 Mar 2002

AB A composition for treating a biofilm structure including a cellular colony and the sessile cells associated with the biofilm structure, comprises an enzyme selected for its ability to dismantle the biofilm structure, an anchor mol. coupled to the enzyme to form an enzyme-anchor complex, the anchor mol. being capable of attaching to a surface on or proximal the biofilm structure, the anchor mol. being selected for its ability to bind to the cellular colony or other bioadhesive mols.. The attachment of the anchor to the surface permits prolonged retention time of the enzyme-anchor complex where the cellular colony and biofilm are present. The first anchor enzyme component degrades biofilm structures and the second anchor enzyme component has the capability to act directly upon the bacteria for a bactericidal effect.

IT 57-92-1, Streptomycin, biological studies

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(comps. for treating **biofilm**)

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD  
(3 CITINGS)

L59 ANSWER 16 OF 30 HCAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2002:221202 HCAPLUS Full-text

DOCUMENT NUMBER: 136:257216

TITLE: Compositions and methods for treating infections using  
cationic peptides alone or in combination with  
antibiotics

INVENTOR(S): Krieger, Timothy J.; Taylor, Robert; Erfle, Douglas;  
Fraser, Janet R.; West, Michael H. P.; McNichol,  
Patricia J.

PATENT ASSIGNEE(S): Micrologix Biotech, Inc, Can.

SOURCE: U.S. Pat. Appl. Publ., 111 pp., Cont.-in-part of U. S.  
6,180,604.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
US 20020035061	A1	20020321	US 1998-30619	19980225
US 6503881	B2	20030107		
US 6180604	B1	20010130	US 1997-915314	19970820
EP 1174439	A2	20020123	EP 2001-119148	19970821
EP 1174439	A3	20030326		
EP 1174439	B1	20081008		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
CA 2282807	A1	19980917	CA 1998-2282807	19980310
AU 9866047	A	19980929	AU 1998-66047	19980310
EP 966481	A2	19991229	EP 1998-907779	19980310
EP 966481	B1	20060719		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 2002544759	T	20021224	JP 1998-538997	19980310
AT 333464	T	20060815	AT 1998-907779	19980310
ES 2264198	T3	20061216	ES 1998-907779	19980310
HK 1025103	A1	20060929	HK 2000-103705	20000620
US 6538106	B1	20030325	US 2000-667486	20000922
US 20030232750	A1	20031218	US 2002-277233	20021018

10/565,591

US 7309759	B2	20071218		
US 20040009910	A1	20040115	US 2003-351985	20030124
US 7390787	B2	20080624		
JP 2005225857	A	20050825	JP 2004-242925	20040823
JP 4073900	B2	20080409		
US 20080242614	A1	20081002	US 2008-58500	20080328
PRIORITY APPLN. INFO.:			US 1996-24754P	P 19960821
			US 1997-34949P	P 19970113
			US 1997-40649P	P 19970310
			US 1997-915314	A2 19970820
			US 1997-60099P	P 19970926
			EP 1997-941352	A3 19970821
			JP 1998-510994	A3 19970821
			US 1998-30619	A 19980225
			WO 1998-CA190	W 19980310
			US 2000-667486	A1 20000922
			US 2003-351985	A1 20030124

OTHER SOURCE(S): MARPAT 136:257216

ED Entered STN: 22 Mar 2002

AB Compns. and methods for treating infections, especially bacterial infections, are provided. Indolicidin peptide analogs containing at least two basic amino acids are prepared. The analogs are administered as modified peptides, preferably containing photo-oxidized solubilizer.

IT **57-92-1**, Streptomycin, biological studies

RL: ADV (Adverse effect, including toxicity); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (compns. and methods for treating infections using cationic peptides alone or in combination with antibiotics)

OS.CITING REF COUNT: 12 THERE ARE 12 CAPLUS RECORDS THAT CITE THIS RECORD (12 CITINGS)

**Text search results: commercial files**

=&gt; d L59 17-30 ibib ab hit

L59 ANSWER 17 OF 30 MEDLINE on STN DUPLICATE 1  
 ACCESSION NUMBER: 2008047574 MEDLINE Full-text  
 DOCUMENT NUMBER: PubMed ID: 17993539  
 TITLE: **Vibrio** parahaemolyticus ScrC modulates cyclic dimeric GMP regulation of gene expression relevant to growth on surfaces.  
 AUTHOR: Ferreira Rosana B R; Antunes Luis Caetano M; Greenberg E Peter; McCarter Linda L  
 CORPORATE SOURCE: Department of Microbiology, The University of Iowa, Iowa City, Iowa 52242, USA.  
 SOURCE: Journal of bacteriology, (2008 Feb) Vol. 190, No. 3, pp. 851-60. Electronic Publication: 2007-11-09. Journal code: 2985120R. E-ISSN: 1098-5530. Report No.: NLM-PMC2223563.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE) (RESEARCH SUPPORT, U.S. GOV'T, NON-P.H.S.)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200802  
 ENTRY DATE: Entered STN: 19 Jan 2008  
 Last Updated on STN: 13 Feb 2008  
 Entered Medline: 12 Feb 2008

AB In **Vibrio** parahaemolyticus, scrC participates in controlling the decision to be a highly mobile swarmer cell or a more adhesive, **biofilm**-proficient cell type. scrC mutants display decreased swarming motility over surfaces and enhanced capsular polysaccharide production. ScrC is a cytoplasmic **membrane** protein that contains both GGDEF and EAL conserved protein domains. These domains have been shown in many organisms to respectively control the formation and degradation of the small signaling nucleotide cyclic dimeric GMP ( **c-di-GMP**). The scrC gene is part of the three-gene scrABC operon. Here we report that this operon influences the cellular nucleotide pool and that **c-di-GMP** levels inversely modulate lateral flagellar and capsular polysaccharide gene expression. High concentrations of this nucleotide prevent swarming and promote adhesiveness. Further, we demonstrate that ScrC has intrinsic diguanylate cyclase and phosphodiesterase activities, and these activities are controlled by ScrAB. Specifically, ScrC acts to form **c-di-GMP** in the absence of ScrA and ScrB; whereas ScrC acts to degrade **c-di-GMP** in the presence of ScrA and ScrB. The scrABC operon is specifically induced by growth on a surface, and the analysis of mutant phenotypes supports a model in which the phosphodiesterase activity of ScrC plays a dominant role during surface translocation and in **biofilms**.

TI **Vibrio** parahaemolyticus ScrC modulates cyclic dimeric GMP regulation of gene expression relevant to growth on surfaces.

AB In **Vibrio** parahaemolyticus, scrC participates in controlling the decision to be a highly mobile swarmer cell or a more adhesive, **biofilm**-proficient cell type. scrC mutants display decreased swarming motility over surfaces and enhanced capsular polysaccharide production. ScrC is a cytoplasmic **membrane** protein that contains both GGDEF and EAL conserved protein domains. These domains have been shown in many organisms to respectively control the formation and degradation of the small signaling nucleotide cyclic dimeric GMP ( **c-di-GMP**). The scrC gene is part of the three-gene scrABC operon. Here we report that this operon influences the cellular nucleotide pool and that **c-di-GMP** levels inversely modulate lateral flagellar and capsular polysaccharide gene expression. High concentrations of this nucleotide prevent swarming and

promote adhesiveness. Further, we demonstrate that ScrC has intrinsic diguanylate cyclase and phosphodiesterase activities, and these activities are controlled by ScrAB. Specifically, ScrC acts to form **c-di-GMP** in the absence of ScrA and ScrB; whereas ScrC acts to degrade **c-di-GMP** in the presence of ScrA and ScrB. The scrABC operon is specifically induced by growth on a surface, and the analysis of mutant phenotypes supports a model in which the phosphodiesterase activity of ScrC plays a dominant role during surface translocation and in **biofilms**.

CT Bacterial Adhesion  
 Bacterial Capsules: GE, genetics  
 Bacterial Capsules: ME, metabolism  
 Bacterial Proteins: CH, chemistry  
 Bacterial Proteins: GE, genetics  
 \*Bacterial Proteins: ME, metabolism  
     **Cyclic GMP: ME, metabolism**  
     **\*Cyclic GMP: PD, pharmacology**  
 Dimerization  
 Flagella: GE, genetics  
 Flagella: ME, metabolism  
 \*Gene Expression Regulation, Bacterial  
     **Membrane Proteins: CH, chemistry**  
     **Membrane Proteins: GE, genetics**  
     **\*Membrane Proteins: ME, metabolism**  
 Mutation  
 Operon  
 Phosphoric Diester Hydrolases: CH, chemistry  
 Phosphoric Diester Hydrolases: GE, genetics  
 Phosphoric Diester Hydrolases: ME, metabolism  
 Phosphorus-Oxygen Lyases: CH, chemistry  
 Phosphorus-Oxygen Lyases: GE, genetics  
 Phosphorus-Oxygen Lyases: ME, metabolism  
     **Vibrio parahaemolyticus: GE, genetics**  
     **\*Vibrio parahaemolyticus: GD, growth & development**  
     **Vibrio parahaemolyticus: ME, metabolism**  
 RN 7665-99-8 (Cyclic GMP)  
 CN 0 (Bacterial Capsules); 0 (Bacterial Proteins); 0 (Membrane  
 Proteins); EC 3.1.4.- (Phosphoric Diester Hydrolases); EC 4.6.-  
 (Phosphorus-Oxygen Lyases); EC 4.6.1.- (diguanylate cyclase)

L59 ANSWER 18 OF 30 MEDLINE on STN DUPLICATE 2  
 ACCESSION NUMBER: 2007303807 MEDLINE Full-text  
 DOCUMENT NUMBER: PubMed ID: 17474125  
 TITLE: Development of nitric oxide synthase-defined neurons in the  
         sea urchin larval ciliary band and evidence for a  
         chemosensory function during metamorphosis.  
 AUTHOR: Bishop Cory D; Brandhorst Bruce P  
 CORPORATE SOURCE: Kewalo Marine Laboratories, University of Hawaii, Honolulu,  
                     Hawaii, USA.. cdbishop@dal.ca  
 SOURCE: Developmental dynamics : an official publication of the  
         American Association of Anatomists, (2007 Jun) Vol. 236,  
         No. 6, pp. 1535-46.  
         Journal code: 9201927. ISSN: 1058-8388.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
                     (RESEARCH SUPPORT, NON-U.S. GOV'T)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200708  
 ENTRY DATE: Entered STN: 23 May 2007  
               Last Updated on STN: 25 Aug 2007

Entered Medline: 24 Aug 2007

- AB We previously reported that initiation of metamorphosis of larvae of *Lytechinus pictus* is negatively regulated by nitric oxide (NO) and **cGMP**. We have examined the expression of nitric oxide synthase (NOS) and **cGMP** in cells of the developing larva. A section of the post-oral ciliary band of feeding larvae includes neural cells defined by their expression of both NOS and the echinoderm neural-specific antibody 1E11. These neurons project processes to the pre-oral neuropile during larval development. Larvae regenerated this section of the ciliary band after its excision, complete with NOS-defined neurons that projected again to the pre-oral neuropile. Excision of ectoderm containing the post-oral ciliary band prevented a behavioral and morphogenetic response of competent larvae to **biofilm**, and **delayed** initiation of metamorphosis. Elevated **cGMP** levels were detected in several larval and juvenile cell types prior to metamorphosis. **Treatment** of larvae with ODQ, an inhibitor of soluble guanylate cyclase, decreased **cGMP** levels and induced metamorphosis while a generator of NO counteracted this effect, indicating inhibition of metamorphosis by NO operates via interaction with soluble guanylate cyclase. We discuss these observations, proposing that the NOS-defined neurons in the post-oral ciliary band have a chemosensory function during settlement and metamorphosis that involves morphologically specialized ectoderm and manipulation of fluid flow. We provide a tentative cellular model of how environmental signals may be transduced into a metamorphic response. Copyright 2007 Wiley-Liss, Inc.
- AB We previously reported that initiation of metamorphosis of larvae of *Lytechinus pictus* is negatively regulated by nitric oxide (NO) and **cGMP**. We have examined the expression of nitric oxide synthase (NOS) and **cGMP** in cells of the developing larva. A section of the post-oral ciliary band of feeding larvae includes neural cells defined by their expression of both NOS and the echinoderm neural-specific antibody 1E11. These neurons project processes to the pre-oral neuropile during larval development. Larvae regenerated this section of the ciliary band after its excision, complete with NOS-defined neurons that projected again to the pre-oral neuropile. Excision of ectoderm containing the post-oral ciliary band prevented a behavioral and morphogenetic response of competent larvae to **biofilm**, and **delayed** initiation of metamorphosis. Elevated **cGMP** levels were detected in several larval and juvenile cell types prior to metamorphosis. **Treatment** of larvae with ODQ, an inhibitor of soluble guanylate cyclase, decreased **cGMP** levels and induced metamorphosis while a generator of NO counteracted this effect, indicating inhibition of metamorphosis by NO operates via interaction with soluble guanylate cyclase. We discuss these observations, proposing that the NOS-defined neurons in the post-oral ciliary band have a chemosensory function during settlement and metamorphosis that involves morphologically specialized ectoderm and manipulation of fluid flow. We provide a tentative cellular model of how environmental signals may be transduced into a metamorphic response. Copyright 2007 Wiley-Liss, Inc.
- CT Aging: PH, physiology  
Animals  
**Biofilms**  
\*Chemotaxis  
**Cyclic GMP: ME, metabolism**  
Gene Expression Regulation, Developmental  
Larva: CY, cytology  
Larva: EN, enzymology  
Larva: GD, growth & development  
\*Metamorphosis, Biological  
Mouth: EN, enzymology  
Mouth: GD, growth & development  
\*Neurons: CY, cytology  
\*Neurons: EN, enzymology  
Nitric Oxide Synthase: GE, genetics



\*Nitric Oxide Synthase: ME, metabolism  
 Sea Urchins: CY, cytology  
 \*Sea Urchins: EN, enzymology  
 Sea Urchins: GE, genetics  
 \*Sea Urchins: GD, growth & development  
 Signal Transduction

RN 7665-99-8 (Cyclic GMP)

L59 ANSWER 19 OF 30 MEDLINE on STN DUPLICATE 3  
 ACCESSION NUMBER: 2008051556 MEDLINE Full-text  
 DOCUMENT NUMBER: PubMed ID: 18028314  
 TITLE: Subcellular location characteristics of the Pseudomonas aeruginosa GGDEF protein, WspR, indicate that it produces cyclic-di-GMP in response to growth on surfaces.  
 AUTHOR: Guvener Zehra Tuzun; Harwood Caroline S  
 CORPORATE SOURCE: Department of Microbiology, University of Washington, Seattle, WA 98195, USA.  
 CONTRACT NUMBER: GM56665 (United States NIGMS NIH HHS)  
 SOURCE: Molecular microbiology, (2007 Dec) Vol. 66, No. 6, pp. 1459-73. Electronic Publication: 2007-11-19.  
 Journal code: 8712028. ISSN: 0950-382X.  
 PUB. COUNTRY: England: United Kingdom  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 (RESEARCH SUPPORT, N.I.H., EXTRAMURAL)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200806  
 ENTRY DATE: Entered STN: 23 Jan 2008  
 Last Updated on STN: 14 Jun 2008  
 Entered Medline: 13 Jun 2008

AB The Pseudomonas aeruginosa Wsp signal transduction system produces cyclic-di-GMP (**c-di-GMP**), an intracellular messenger that stimulates **biofilm** formation and suppresses motility. The Wsp system is homologous to chemotaxis systems and includes a **membrane**-bound receptor protein, WspA, and a response regulator GGDEF protein, WspR, that catalyses **c-di-GMP** synthesis when phosphorylated. We found that the subcellular distributions of fluorescent protein-tagged WspA and WspR differed markedly from their chemotaxis counterparts. WspA-YFP formed patches in cells whereas WspR-YFP was dispersed when unphosphorylated and formed bright cytoplasmic clusters when phosphorylated. WspR formed clusters in cells of a DeltawspF mutant, a genetic background that causes constitutive phosphorylation of WspR, but was dispersed in cells of a wspA mutant, a genetic background necessary for WspR phosphorylation. In addition, WspR mutated at Asp70, its predicted site of phosphorylation, did not form clusters. **C-di-GMP** synthesis was not required for cluster formation. WspR-YFP was dispersed in liquid-grown wild-type cells, but formed clusters that sometimes appeared and disappeared over the course of a few minutes in cells grown on an agar surface. Our results suggest that the compartmentalized production of **c-di-GMP** in response to a stimulus associated with growth on a surface is an important functional characteristic of the Wsp system.

AB The Pseudomonas aeruginosa Wsp signal transduction system produces cyclic-di-GMP (**c-di-GMP**), an intracellular messenger that stimulates **biofilm** formation and suppresses motility. The Wsp system is homologous to chemotaxis systems and includes a **membrane**-bound receptor protein, WspA, and a response regulator GGDEF protein, WspR, that catalyses **c-di-GMP** synthesis when phosphorylated. We found that the subcellular distributions of fluorescent protein-tagged WspA and WspR differed markedly from their chemotaxis counterparts. WspA-YFP formed patches in cells whereas WspR-YFP was dispersed when unphosphorylated and formed bright cytoplasmic clusters when phosphorylated. WspR formed clusters in cells of a DeltawspF mutant, a genetic background that causes constitutive phosphorylation of WspR, but was dispersed in cells of a wspA

mutant, a genetic background necessary for WspR phosphorylation. In addition, WspR mutated at Asp70, its predicted site of phosphorylation, did not form clusters. **C-di-GMP** synthesis was not required for cluster formation. WspR-YFP was dispersed in liquid-grown wild-type cells, but formed clusters that sometimes appeared and disappeared over the course of a few minutes in cells grown on an agar surface. Our results suggest that the compartmentalized production of **c-di-GMP** in response to a stimulus associated with growth on a surface is an important functional characteristic of the Wsp system.

CT Bacterial Proteins: GE, genetics

\*Bacterial Proteins: ME, metabolism

\*Cyclic GMP: AA, analogs & derivatives

Cyclic GMP: ME, metabolism

Cytoplasm: ME, metabolism

Microscopy, Fluorescence

Models, Biological

Mutagenesis, Site-Directed

Mutation

Phosphorylation

Pseudomonas aeruginosa: GE, genetics

\*Pseudomonas aeruginosa: ME, metabolism

Recombinant Fusion Proteins: GE, genetics

\*Recombinant Fusion Proteins: ME, metabolism

Signal Transduction: GE, genetics

Signal Transduction: PH, physiology

RN 61093-23-0 (bis(3',5')-cyclic diguanylic acid); 7665-99-8  
(Cyclic GMP)

L59 ANSWER 20 OF 30 MEDLINE on STN DUPLICATE 4  
 ACCESSION NUMBER: 2006671047 MEDLINE Full-text  
 DOCUMENT NUMBER: PubMed ID: 16980460  
 TITLE: Diguanylate cyclases control magnesium-dependent motility  
 of **Vibrio** fischeri.  
 AUTHOR: O'Shea Therese M; Klein Adam H; Geszvain Kati; Wolfe Alan  
 J; Visick Karen L  
 CORPORATE SOURCE: Department of Microbiology and Immunology, Loyola  
 University Chicago, 2160 S. First Ave., Bldg. 105, Maywood,  
 IL 60153, USA.  
 CONTRACT NUMBER: GM066130 (United States NIGMS NIH HHS)  
 GM59690 (United States NIGMS NIH HHS)  
 SOURCE: Journal of bacteriology, (2006 Dec) Vol. 188, No. 23, pp.  
 8196-205. Electronic Publication: 2006-09-15.  
 Journal code: 2985120R. ISSN: 0021-9193.  
 Report No.: NLM-PMC1698204.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 (RESEARCH SUPPORT, N.I.H., EXTRAMURAL)  
 (RESEARCH SUPPORT, NON-U.S. GOV'T)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200612  
 ENTRY DATE: Entered STN: 21 Nov 2006  
 Last Updated on STN: 28 Dec 2006  
 Entered Medline: 27 Dec 2006

AB Flagellar biogenesis and hence motility of **Vibrio** fischeri depends upon the presence of magnesium. In the absence of magnesium, cells contain few or no flagella and are poorly motile or nonmotile. To dissect the mechanism by which this regulation occurs, we screened transposon insertion mutants for those that could migrate through soft agar medium lacking added magnesium. We identified mutants with insertions in two distinct genes, VF0989 and VFA0959, which we termed mifA and mifB, respectively, for magnesium-dependent induction

of flagellation. Each gene encodes a predicted **membrane**-associated protein with diguanylate cyclase activity. Consistent with that activity, introduction into *V. fischeri* of medium-copy plasmids carrying these genes inhibited motility. Furthermore, multicopy expression of *mifA* induced other phenotypes known to be correlated with diguanylate cyclase activity, including cellulose biosynthesis and **biofilm** formation. To directly test their function, we introduced the wild-type genes on high-copy plasmids into *Escherichia coli*. We assayed for the production of cyclic di-GMP using two-dimensional thin-layer chromatography and found that strains carrying these plasmids produced a small but reproducible spot that migrated with an R(f) value consistent with cyclic di-GMP that was not produced by strains carrying the vector control. Disruptions of *mifA* or *mifB* increased flagellin levels, while multicopy expression decreased them. Semiquantitative reverse transcription-PCR experiments revealed no significant difference in the amount of flagellin transcripts produced in either the presence or absence of Mg(2+) by either vector control or *mifA*-overexpressing cells, indicating that the impact of magnesium and cyclic-di-GMP primarily acts following transcription. Finally, we present a model for the roles of magnesium and cyclic di-GMP in the control of motility of *V. fischeri*.

TI Diguanylate cyclases control magnesium-dependent motility of **Vibrio fischeri**.

AB Flagellar biogenesis and hence motility of **Vibrio fischeri** depends upon the presence of magnesium. In the absence of magnesium, cells contain few or no flagella and are poorly motile or nonmotile. To dissect the mechanism by which this regulation occurs, we screened transposon insertion mutants for those that could migrate through soft agar medium lacking added magnesium. We identified mutants with insertions in two distinct genes, VF0989 and VFA0959, which we termed *mifA* and *mifB*, respectively, for magnesium-dependent induction of flagellation. Each gene encodes a predicted **membrane**-associated protein with diguanylate cyclase activity. Consistent with that activity, introduction into *V. fischeri* of medium-copy plasmids carrying these genes inhibited motility. Furthermore, multicopy expression of *mifA* induced other phenotypes known to be correlated with diguanylate cyclase activity, including cellulose biosynthesis and **biofilm** formation. To directly test their function, we introduced the wild-type genes on high-copy plasmids into *Escherichia coli*. We assayed for the production of cyclic di-GMP using two-dimensional thin-layer chromatography and found that strains carrying these plasmids produced a small but reproducible spot that migrated with an R(f) value consistent with cyclic di-GMP that was not produced by strains carrying the vector control. Disruptions of *mifA* or *mifB* increased flagellin levels, while multicopy expression decreased them. Semiquantitative reverse transcription-PCR experiments revealed no significant difference in the amount of flagellin transcripts produced in either the presence or absence of Mg(2+) by either vector control or *mifA*-overexpressing cells, indicating that the impact of magnesium and cyclic-di-GMP primarily acts following transcription. Finally, we present a model for the roles of magnesium and cyclic di-GMP in the control of motility of *V. fischeri*.

CT Bacterial Proteins: GE, genetics

\*Bacterial Proteins: PH, physiology

**Biofilms: DE, drug effects**

Cellulose: BI, biosynthesis

**Cyclic GMP: PH, physiology**

\*Down-Regulation

*Escherichia coli*: GE, genetics

*Escherichia coli*: ME, metabolism

Flagella

\*Gene Expression Regulation, Bacterial

Genetic Vectors

Locomotion

\*Magnesium: PH, physiology

Mutagenesis, Insertional  
 Phosphorus-Oxygen Lyases: GE, genetics  
 \*Phosphorus-Oxygen Lyases: PH, physiology  
 Plasmids  
 Transfection

**Vibrio fischeri: GE, genetics**  
**\*Vibrio fischeri: PH, physiology**

RN 7439-95-4 (Magnesium); 7665-99-8 (Cyclic GMP); 9004-34-6  
 (Cellulose)

L59 ANSWER 21 OF 30 MEDLINE on STN

ACCESSION NUMBER: 2009181697 MEDLINE Full-text

DOCUMENT NUMBER: PubMed ID: 19218451

TITLE: LapD is a bis-(3',5')-cyclic dimeric GMP-binding protein  
 that regulates surface attachment by Pseudomonas  
 fluorescens Pf0-1.

AUTHOR: Newell Peter D; Monds Russell D; O'Toole George A

CORPORATE SOURCE: Department of Microbiology and Immunology, Dartmouth  
 Medical School, Hanover, NH 03755, USA.

CONTRACT NUMBER: T32 GM08704 (United States NIGMS NIH HHS)

SOURCE: Proceedings of the National Academy of Sciences of the  
 United States of America, (2009 Mar 3) Vol. 106, No. 9, pp.  
 3461-6. Electronic Publication: 2009-02-13.  
 Journal code: 7505876. E-ISSN: 1091-6490.  
 Report No.: NLM-PMC2651287.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 (RESEARCH SUPPORT, N.I.H., EXTRAMURAL)  
 (RESEARCH SUPPORT, U.S. GOV'T, NON-P.H.S.)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200904

ENTRY DATE: Entered STN: 5 Mar 2009  
 Last Updated on STN: 2 Apr 2009  
 Entered Medline: 1 Apr 2009

AB The second messenger cyclic dimeric GMP (**c-di- GMP**) regulates surface  
 attachment and **biofilm** formation by many bacteria. For Pseudomonas  
 fluorescens Pf0-1, **c- di-GMP** impacts the secretion and localization of the  
 adhesin LapA, which is absolutely required for stable surface attachment and  
**biofilm** formation by this bacterium. In this study we characterize LapD, a  
 unique **c-di-GMP** effector protein that **controls biofilm** formation by  
 communicating intracellular **c-di-GMP** levels to the **membrane**-localized  
 attachment machinery via its periplasmic domain. LapD contains degenerate and  
 enzymatically inactive diguanylate cyclase and **c-di-GMP** phosphodiesterase  
 (EAL) domains and binds to **c-di- GMP** through a degenerate EAL domain. We  
 present evidence that LapD utilizes an inside-out signaling mechanism: binding  
**c- di-GMP** in the cytoplasm and communicating this signal to the periplasm via  
 its periplasmic domain. Furthermore, we show that LapD serves as the **c-di-GMP**  
 receptor connecting environmental modulation of intracellular **c-di- GMP** levels  
 by inorganic phosphate to regulation of LapA localization and thus surface  
 commitment by P. fluorescens.

AB The second messenger cyclic dimeric GMP (**c-di- GMP**) regulates surface  
 attachment and **biofilm** formation by many bacteria. For Pseudomonas  
 fluorescens Pf0-1, **c- di-GMP** impacts the secretion and localization of the  
 adhesin LapA, which is absolutely required for stable surface attachment and  
**biofilm** formation by this bacterium. In this study we characterize LapD, a  
 unique **c-di-GMP** effector protein that **controls biofilm** formation by  
 communicating intracellular **c-di-GMP** levels to the **membrane**-localized  
 attachment machinery via its periplasmic domain. LapD contains degenerate and  
 enzymatically inactive diguanylate cyclase and **c-di-GMP** phosphodiesterase

(EAL) domains and binds to **c-di-GMP** through a degenerate EAL domain. We present evidence that LapD utilizes an inside-out signaling mechanism: binding **c-di-GMP** in the cytoplasm and communicating this signal to the periplasm via its periplasmic domain. Furthermore, we show that LapD serves as the **c-di-GMP** receptor connecting environmental modulation of intracellular **c-di-GMP** levels by inorganic phosphate to regulation of LapA localization and thus surface commitment by *P. fluorescens*.

CT **3',5'-Cyclic-GMP Phosphodiesterases: ME, metabolism**

Adhesins, Bacterial: ME, metabolism

\*Bacterial Adhesion

Carrier Proteins: GE, genetics

\*Carrier Proteins: ME, metabolism

**Cyclic GMP: AA, analogs & derivatives**

**Cyclic GMP: ME, metabolism**

Enzyme Activation

Intracellular Signaling Peptides and Proteins: GE, genetics

\*Intracellular Signaling Peptides and Proteins: ME, metabolism

Mutation: GE, genetics

Protein Binding

\*Protein Multimerization

*Pseudomonas fluorescens*: GE, genetics

\**Pseudomonas fluorescens*: ME, metabolism

Signal Transduction

RN **61093-23-0 (bis(3',5')-cyclic diguanylic acid); 7665-99-8 (Cyclic GMP)**

CN 0 (Adhesins, Bacterial); 0 (Carrier Proteins); 0 (Intracellular Signaling Peptides and Proteins); 0 (**cyclic GMP-binding protein**); EC 3.1.4.35 (**3',5'-Cyclic-GMP Phosphodiesterases**)

L59 ANSWER 22 OF 30 MEDLINE on STN

ACCESSION NUMBER: 2009444210 MEDLINE Full-text

DOCUMENT NUMBER: PubMed ID: 19460094

TITLE: Second messenger signalling governs *Escherichia coli* **biofilm** induction upon ribosomal stress.

AUTHOR: Boehm Alex; Steiner Samuel; Zaehring Franziska; Casanova Alain; Hamburger Fabienne; Ritz Daniel; Keck Wolfgang; Ackermann Martin; Schirmer Tilman; Jenal Urs

CORPORATE SOURCE: Biozentrum, University of Basel, Klingelbergstrasse 50/70, 4056 Basel, Switzerland.. alexander.boehm@unibas.ch

SOURCE: Molecular microbiology, (2009 Jun) Vol. 72, No. 6, pp. 1500-16. Electronic Publication: 2009-05-15. Journal code: 8712028. E-ISSN: 1365-2958.

PUB. COUNTRY: England: United Kingdom

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
(RESEARCH SUPPORT, NON-U.S. GOV'T)  
(RESEARCH SUPPORT, U.S. GOV'T, NON-P.H.S.)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200907

ENTRY DATE: Entered STN: 27 Jun 2009

Last Updated on STN: 22 Jul 2009

Entered Medline: 21 Jul 2009

AB **Biofilms** are communities of surface-attached, matrix-embedded microbial cells that can resist antimicrobial chemotherapy and contribute to persistent infections. Using an *Escherichia coli* **biofilm** model we found that exposure of bacteria to subinhibitory concentrations of ribosome-targeting antibiotics leads to strong **biofilm** induction. We present evidence that this effect is elicited by the ribosome in response to translational stress. **Biofilm** induction involves upregulation of the polysaccharide adhesin poly-beta-1,6-N-

acetyl-glucosamine (poly-GlcNAc) and two components of the poly-GlcNAc biosynthesis machinery, PgaA and PgaD. Poly-GlcNAc **control** depends on the **bacterial** signalling molecules guanosine-bis 3', 5'(diphosphate) (ppGpp) and bis-(3'-5 ')-**cyclic di-GMP (c-di-GMP)**. **Treatment** with translation inhibitors causes a ppGpp hydrolase (SpoT)-mediated reduction of ppGpp levels, resulting in specific derepression of PgaA. Maximal induction of PgaD and poly-GlcNAc synthesis requires the production of **c-di-GMP** by the dedicated diguanylate cyclase YdeH. Our results identify a novel regulatory mechanism that relies on ppGpp signalling to relay information about ribosomal performance to the Pga machinery, thereby inducing adhesin production and **biofilm** formation. Based on the important synergistic roles of ppGpp and **c-di-GMP** in this process, we suggest that interference with bacterial second messenger signalling might represent an effective means for **biofilm control** during chronic **infections**.

TI Second messenger signalling governs Escherichia coli **biofilm** induction upon ribosomal stress.

AB **Biofilms** are communities of surface-attached, matrix-embedded microbial cells that can resist antimicrobial chemotherapy and contribute to persistent infections. Using an Escherichia coli **biofilm** model we found that exposure of bacteria to subinhibitory concentrations of ribosome-targeting antibiotics leads to strong **biofilm** induction. We present evidence that this effect is elicited by the ribosome in response to translational stress. **Biofilm** induction involves upregulation of the polysaccharide adhesin poly-beta-1,6-N-acetyl-glucosamine (poly-GlcNAc) and two components of the poly-GlcNAc biosynthesis machinery, PgaA and PgaD. Poly-GlcNAc **control** depends on the **bacterial** signalling molecules guanosine-bis 3', 5'(diphosphate) (ppGpp) and bis-(3'-5 ')-**cyclic di-GMP (c-di-GMP)**. **Treatment** with translation inhibitors causes a ppGpp hydrolase (SpoT)-mediated reduction of ppGpp levels, resulting in specific derepression of PgaA. Maximal induction of PgaD and poly-GlcNAc synthesis requires the production of **c-di-GMP** by the dedicated diguanylate cyclase YdeH. Our results identify a novel regulatory mechanism that relies on ppGpp signalling to relay information about ribosomal performance to the Pga machinery, thereby inducing adhesin production and **biofilm** formation. Based on the important synergistic roles of ppGpp and **c-di-GMP** in this process, we suggest that interference with bacterial second messenger signalling might represent an effective means for **biofilm control** during chronic **infections**.

CT Adhesins, Bacterial: ME, metabolism  
 Anti-Bacterial Agents: PD, pharmacology  
 \*Biofilms: GD, growth & development  
 \*Cyclic GMP: AA, analogs & derivatives  
 Cyclic GMP: ME, metabolism  
 Escherichia coli: DE, drug effects  
 Escherichia coli: GE, genetics  
 Escherichia coli: ME, metabolism  
 \*Escherichia coli: PH, physiology  
 Escherichia coli Proteins: GE, genetics  
 Escherichia coli Proteins: ME, metabolism  
 Gene Expression Regulation, Bacterial  
 \*Guanosine Tetraphosphate: ME, metabolism  
 Phosphorus-Oxygen Lyases: GE, genetics  
 Phosphorus-Oxygen Lyases: ME, metabolism  
 Protein Biosynthesis: DE, drug effects  
 Pyrophosphatases: GE, genetics  
 Pyrophosphatases: ME, metabolism  
 RNA Processing, Post-Transcriptional  
 \*Ribosomes: DE, drug effects  
 \*Second Messenger Systems  
 beta-Glucans: ME, metabolism  
 RN 33503-72-9 (Guanosine Tetraphosphate); 61093-23-0 (bis(3',5')-cyclic

## diguanylic acid); 7665-99-8 (Cyclic GMP)

L59 ANSWER 23 OF 30 MEDLINE on STN

ACCESSION NUMBER: 2009124691 MEDLINE Full-text

DOCUMENT NUMBER: PubMed ID: 19088322

TITLE: MucR, a novel **membrane**-associated regulator of alginate biosynthesis in *Pseudomonas aeruginosa*.

AUTHOR: Hay Iain D; Remminghorst Uwe; Rehm Bernd H A

CORPORATE SOURCE: Institute of Molecular Biosciences, Massey University, Private Bag 11222, Palmerston North, New Zealand.

SOURCE: Applied and environmental microbiology, (2009 Feb) Vol. 75, No. 4, pp. 1110-20. Electronic Publication: 2008-12-16. Journal code: 7605801. E-ISSN: 1098-5336.

Report No.: NLM-PMC2643583.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
(RESEARCH SUPPORT, NON-U.S. GOV'T)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200902

ENTRY DATE: Entered STN: 10 Feb 2009

Last Updated on STN: 24 Feb 2009

Entered Medline: 20 Feb 2009

AB Alginate biosynthesis by *Pseudomonas aeruginosa* was shown to be regulated by the intracellular second messenger bis-(3'-5')-cyclic-dimeric-GMP ( **c-di-GMP**), and binding of **c- di-GMP** to the **membrane** protein Alg44 was required for alginate production. In this study, PA1727, a **c- di-GMP**-synthesizing enzyme was functionally analyzed and identified to be involved in regulation of alginate production. Deletion of the PA1727 gene in the mucoid alginate-overproducing *P. aeruginosa* strain PDO300 resulted in a nonmucoid phenotype and an about 38-fold decrease in alginate production; thus, this gene is designated mucR. The mucoid alginate-overproducing phenotype was restored by introducing the mucR gene into the isogenic DeltamucR mutant. Moreover, transfer of the MucR-encoding plasmid into strain PDO300 led to an about sevenfold increase in alginate production, wrinkly colony morphology, increased pellicle formation, auto-aggregation, and the formation of highly structured **biofilms** as well as the **inhibition** of swarming motility. Outer **membrane** protein profile analysis showed that overproduction of MucR mediates a strong reduction in the copy number of FliC (flagellin), required for flagellum-mediated motility. Translational reporter enzyme fusions with LacZ and PhoA suggested that MucR is located in the cytoplasmic **membrane** with a cytosolic C terminus. Deletion of the proposed C-terminal GGDEF domain abolished MucR function. MucR was purified and identified using tryptic peptide fingerprinting and matrix-assisted laser desorption ionization-time of flight mass spectrometry. Overall, experimental evidence was provided suggesting that MucR specifically regulates alginate biosynthesis by activation of alginate production through generation of a localized **c-di-GMP** pool in the vicinity of Alg44.

TI MucR, a novel **membrane**-associated regulator of alginate biosynthesis in *Pseudomonas aeruginosa*.

AB Alginate biosynthesis by *Pseudomonas aeruginosa* was shown to be regulated by the intracellular second messenger bis-(3'-5')-cyclic-dimeric-GMP ( **c-di-GMP**), and binding of **c- di-GMP** to the **membrane** protein Alg44 was required for alginate production. In this study, PA1727, a **c- di-GMP**-synthesizing enzyme was functionally analyzed and identified to be involved in regulation of alginate production. Deletion of the PA1727 gene in the mucoid alginate-overproducing *P. aeruginosa* strain PDO300 resulted in a nonmucoid phenotype and an about 38-fold decrease in alginate production; thus, this gene is designated mucR. The mucoid alginate-overproducing phenotype was restored by introducing the mucR gene into the isogenic DeltamucR mutant. Moreover,

transfer of the MucR-encoding plasmid into strain PDO300 led to an about sevenfold increase in alginate production, wrinkly colony morphology, increased pellicle formation, auto-aggregation, and the formation of highly structured **biofilms** as well as the **inhibition** of swarming motility. Outer **membrane** protein profile analysis showed that overproduction of MucR mediates a strong reduction in the copy number of *FliC* (flagellin), required for flagellum-mediated motility. Translational reporter enzyme fusions with *LacZ* and *PhoA* suggested that MucR is located in the cytoplasmic **membrane** with a cytosolic C terminus. Deletion of the proposed C-terminal GGDEF domain abolished MucR function. MucR was purified and identified using tryptic peptide fingerprinting and matrix-assisted laser desorption ionization-time of flight mass spectrometry. Overall, experimental evidence was provided suggesting that MucR specifically regulates alginate biosynthesis by activation of alginate production through generation of a localized **c-di-GMP** pool in the vicinity of Alg44.

CT Alginates  
 Alkaline Phosphatase: GE, genetics  
 Alkaline Phosphatase: ME, metabolism  
**Bacterial Outer Membrane Proteins: AN, analysis**  
 Bacterial Proteins: GE, genetics  
 Bacterial Proteins: IP, isolation & purification  
 \*Bacterial Proteins: PH, physiology  
**Cell Membrane: CH, chemistry**  
**Cyclic GMP: AA, analogs & derivatives**  
**Cyclic GMP: ME, metabolism**  
 Gene Deletion  
 \*Gene Expression Regulation, Bacterial  
 Genes, Reporter  
 Genetic Complementation Test  
 Glucuronic Acid: BI, biosynthesis  
 Hexuronic Acids  
 Locomotion  
 Pseudomonas aeruginosa: CH, chemistry  
 Pseudomonas aeruginosa: GE, genetics  
 Pseudomonas aeruginosa: ME, metabolism  
 \*Pseudomonas aeruginosa: PH, physiology  
 Recombinant Fusion Proteins: GE, genetics  
 Recombinant Fusion Proteins: ME, metabolism  
 Spectrometry, Mass, Matrix-Assisted Laser Desorption-Ionization  
 beta-Galactosidase: GE, genetics  
 beta-Galactosidase: ME, metabolism  
 RN 576-37-4 (Glucuronic Acid); **61093-23-0 (bis(3',5')-cyclic diguanylic acid); 7665-99-8 (Cyclic GMP)**; 9005-32-7 (alginic acid)  
 CN 0 (Alginates); 0 (Bacterial Outer **Membrane** Proteins); 0 (Bacterial Proteins); 0 (Hexuronic Acids); 0 (Recombinant Fusion Proteins); EC 3.1.3.1 (Alkaline Phosphatase); EC 3.2.1.23 (beta-Galactosidase)

L59 ANSWER 24 OF 30 MEDLINE on STN  
 ACCESSION NUMBER: 2008457940 MEDLINE Full-text  
 DOCUMENT NUMBER: PubMed ID: 18502872  
 TITLE: A **staphylococcal** GGDEF domain protein regulates **biofilm** formation independently of cyclic dimeric GMP.  
 AUTHOR: Holland Linda M; O'Donnell Sinead T; Ryjenkov Dmitri A; Gomelsky Larissa; Slater Shawn R; Fey Paul D; Gomelsky Mark; O'Gara James P  
 CORPORATE SOURCE: School of Biomolecular and Biomedical Science, Ardmore House, University College Dublin, Belfield, Dublin 4, Ireland.



CONTRACT NUMBER: AI49311 (United States NIAID NIH HHS)

SOURCE: Journal of bacteriology, (2008 Aug) Vol. 190, No. 15, pp. 5178-89. Electronic Publication: 2008-05-23.  
Journal code: 2985120R. E-ISSN: 1098-5530.  
Report No.: NLM-PMC2493275.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
(RESEARCH SUPPORT, N.I.H., EXTRAMURAL)  
(RESEARCH SUPPORT, NON-U.S. GOV'T)  
(RESEARCH SUPPORT, U.S. GOV'T, NON-P.H.S.)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200808

ENTRY DATE: Entered STN: 19 Jul 2008  
Last Updated on STN: 13 Aug 2008  
Entered Medline: 12 Aug 2008

AB Cyclic dimeric GMP (**c-di-GMP**) is an important **biofilm** regulator that allosterically activates enzymes of exopolysaccharide biosynthesis. Proteobacterial genomes usually encode multiple GGDEF domain-containing diguanylate cyclases responsible for **c-di-GMP** synthesis. In contrast, only one conserved GGDEF domain protein, GdpS (for GGDEF domain protein from **Staphylococcus**), and a second protein with a highly modified GGDEF domain, GdpP, are present in the sequenced **staphylococcal** genomes. Here, we investigated the role of GdpS in **biofilm** formation in **Staphylococcus epidermidis**. Inactivation of gdpS impaired **biofilm** formation in medium supplemented with NaCl under static and flow-cell conditions, whereas gdpS overexpression complemented the mutation and enhanced wild-type **biofilm** development. GdpS increased production of the icaADBC-encoded exopolysaccharide, poly-N-acetyl-glucosamine, by elevating icaADBC mRNA levels. Unexpectedly, **c-di-GMP** synthesis was found to be irrelevant for the ability of GdpS to elevate icaADBC expression. Mutagenesis of the GGDEF motif essential for diguanylate cyclase activity did not impair GdpS, and the N-terminal fragment of GdpS lacking the GGDEF domain partially complemented the gdpS mutation. Furthermore, heterologous diguanylate cyclases expressed in trans failed to complement the gdpS mutation, and the purified GGDEF domain from GdpS possessed no diguanylate cyclase activity in vitro. The gdpS gene from **Staphylococcus aureus** exhibited similar characteristics to its **S. epidermidis** ortholog, suggesting that the GdpS-mediated signal transduction is conserved in **staphylococci**. Therefore, GdpS affects **biofilm** formation through a novel **c-di-GMP**-independent mechanism involving increased icaADBC mRNA levels and exopolysaccharide biosynthesis. Our data raise the possibility that **staphylococci** cannot synthesize **c-di-GMP** and have only remnants of a **c-di-GMP** signaling pathway.

TI A **staphylococcal** GGDEF domain protein regulates **biofilm** formation independently of cyclic dimeric GMP.

AB Cyclic dimeric GMP (**c-di-GMP**) is an important **biofilm** regulator that allosterically activates enzymes of exopolysaccharide biosynthesis. Proteobacterial genomes usually encode multiple GGDEF domain-containing diguanylate cyclases responsible for **c-di-GMP** synthesis. In contrast, only one conserved GGDEF domain protein, GdpS (for GGDEF domain protein from **Staphylococcus**), and a second protein with a highly modified GGDEF domain, GdpP, are present in the sequenced **staphylococcal** genomes. Here, we investigated the role of GdpS in **biofilm** formation in **Staphylococcus epidermidis**. Inactivation of gdpS impaired **biofilm** formation in medium supplemented with NaCl under static and flow-cell conditions, whereas gdpS overexpression complemented the mutation and enhanced wild-type **biofilm** development. GdpS increased production of the icaADBC-encoded exopolysaccharide, poly-N-acetyl-glucosamine, by elevating icaADBC mRNA levels. Unexpectedly, **c-di-GMP** synthesis was found to be irrelevant for the ability of GdpS to elevate icaADBC expression. Mutagenesis of the GGDEF motif

essential for diguanylate cyclase activity did not impair GdpS, and the N-terminal fragment of GdpS lacking the GGDEF domain partially complemented the gdpS mutation. Furthermore, heterologous diguanylate cyclases expressed in trans failed to complement the gdpS mutation, and the purified GGDEF domain from GdpS possessed no diguanylate cyclase activity in vitro. The gdpS gene from *Staphylococcus aureus* exhibited similar characteristics to its *S. epidermidis* ortholog, suggesting that the GdpS-mediated signal transduction is conserved in *staphylococci*. Therefore, GdpS affects **biofilm** formation through a novel **c-di-GMP** -independent mechanism involving increased icaADBC mRNA levels and exopolysaccharide biosynthesis. Our data raise the possibility that *staphylococci* cannot synthesize **c-di-GMP** and have only remnants of a **c-di-GMP** signaling pathway.

CT Amino Acid Sequence

\*Biofilms: GD, growth & development

\*Cyclic GMP: AA, analogs & derivatives

Cyclic GMP: ME, metabolism

Gene Deletion

Gene Dosage

Gene Expression Profiling

\*Gene Expression Regulation

Genetic Complementation Test

Molecular Sequence Data

Mutagenesis, Insertional

Mutagenesis, Site-Directed

Mutation

Phosphorus-Oxygen Lyases: GE, genetics

\*Phosphorus-Oxygen Lyases: ME, metabolism

Polysaccharides, Bacterial: BI, biosynthesis

Sequence Alignment

Sequence Deletion

*Staphylococcus aureus*: EN, enzymology

*Staphylococcus aureus*: GE, genetics

*Staphylococcus epidermidis*: EN, enzymology

*Staphylococcus epidermidis*: GE, genetics

\**Staphylococcus epidermidis*: PH, physiology

RN 61093-23-0 (bis(3',5')-cyclic diguanylic acid); 7665-99-8  
(Cyclic GMP)

L59 ANSWER 25 OF 30 MEDLINE on STN

ACCESSION NUMBER: 2007652249 MEDLINE [Full-text](#)

DOCUMENT NUMBER: PubMed ID: 17586641

TITLE: BifA, a cyclic-Di-GMP phosphodiesterase, inversely regulates **biofilm** formation and swarming motility by *Pseudomonas aeruginosa* PA14.

AUTHOR: Kuchma Sherry L; Brothers Kimberly M; Merritt Judith H; Liberati Nicole T; Ausubel Frederick M; O'Toole George A

CORPORATE SOURCE: Department of Microbiology and Immunology, Dartmouth Medical School, Rm. 505, Vail Building, North College St., Hanover, NH 03755, USA.

CONTRACT NUMBER: 1-P20-RR01878 (United States NCRR NIH HHS)  
AI51360 (United States NIAID NIH HHS)

SOURCE: Journal of bacteriology, (2007 Nov) Vol. 189, No. 22, pp. 8165-78. Electronic Publication: 2007-06-22.  
Journal code: 2985120R. E-ISSN: 1098-5530.  
Report No.: NLM-PMC2168662.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
(RESEARCH SUPPORT, N.I.H., EXTRAMURAL)  
(RESEARCH SUPPORT, NON-U.S. GOV'T)  
(RESEARCH SUPPORT, U.S. GOV'T, NON-P.H.S.)

LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200801  
 ENTRY DATE: Entered STN: 6 Nov 2007  
 Last Updated on STN: 15 Jan 2008  
 Entered Medline: 14 Jan 2008

AB The intracellular signaling molecule, cyclic-di-GMP (**c-di-GMP**), has been shown to influence bacterial behaviors, including motility and **biofilm** formation. We report the identification and characterization of PA4367, a gene involved in regulating surface-associated behaviors in *Pseudomonas aeruginosa*. The PA4367 gene encodes a protein with an EAL domain, associated with **c-di-GMP** phosphodiesterase activity, as well as a GGDEF domain, which is associated with a **c-di-GMP**-synthesizing diguanylate cyclase activity. Deletion of the PA4367 gene results in a severe defect in swarming motility and a hyperbiofilm phenotype; thus, we designate this gene *bifA*, for **biofilm** formation. We show that BifA localizes to the inner **membrane** and, in biochemical studies, that purified BifA protein exhibits phosphodiesterase activity in vitro but no detectable diguanylate cyclase activity. Furthermore, mutational analyses of the conserved EAL and GGDEF residues of BifA suggest that both domains are important for the observed phosphodiesterase activity. Consistent with these data, the *DeltabifA* mutant exhibits increased cellular pools of **c-di-GMP** relative to the wild type and increased synthesis of a polysaccharide produced by the *pel* locus. This increased polysaccharide production is required for the enhanced **biofilm** formed by the *DeltabifA* mutant but does not contribute to the observed swarming defect. The *DeltabifA* mutation also results in decreased flagellar reversals. Based on epistasis studies with the previously described *sadB* gene, we propose that BifA functions upstream of SadB in the **control** of **biofilm** formation and swarming.

TI *BifA*, a cyclic-Di-GMP phosphodiesterase, inversely regulates **biofilm** formation and swarming motility by *Pseudomonas aeruginosa* PA14.

AB The intracellular signaling molecule, cyclic-di-GMP (**c-di-GMP**), has been shown to influence bacterial behaviors, including motility and **biofilm** formation. We report the identification and characterization of PA4367, a gene involved in regulating surface-associated behaviors in *Pseudomonas aeruginosa*. The PA4367 gene encodes a protein with an EAL domain, associated with **c-di-GMP** phosphodiesterase activity, as well as a GGDEF domain, which is associated with a **c-di-GMP**-synthesizing diguanylate cyclase activity. Deletion of the PA4367 gene results in a severe defect in swarming motility and a hyperbiofilm phenotype; thus, we designate this gene *bifA*, for **biofilm** formation. We show that BifA localizes to the inner **membrane** and, in biochemical studies, that purified BifA protein exhibits phosphodiesterase activity in vitro but no detectable diguanylate cyclase activity. Furthermore, mutational analyses of the conserved EAL and GGDEF residues of BifA suggest that both domains are important for the observed phosphodiesterase activity. Consistent with these data, the *DeltabifA* mutant exhibits increased cellular pools of **c-di-GMP** relative to the wild type and increased synthesis of a polysaccharide produced by the *pel* locus. This increased polysaccharide production is required for the enhanced **biofilm** formed by the *DeltabifA* mutant but does not contribute to the observed swarming defect. The *DeltabifA* mutation also results in decreased flagellar reversals. Based on epistasis studies with the previously described *sadB* gene, we propose that BifA functions upstream of SadB in the **control** of **biofilm** formation and swarming.

CT Bacterial Proteins: GE, genetics  
 Bacterial Proteins: ME, metabolism  
 \*Biofilms: GD, growth & development  
 Cell Membrane  
 \*Cyclic GMP: AA, analogs & derivatives

**Cyclic GMP: ME, metabolism**Gene Expression Regulation, Bacterial  
Movement

Phosphoric Diester Hydrolases: GE, genetics

\*Phosphoric Diester Hydrolases: ME, metabolism

Protein Transport

\*Pseudomonas aeruginosa: CY, cytology

\*Pseudomonas aeruginosa: EN, enzymology

RN **61093-23-0 (bis(3',5')-cyclic diguanylic acid); 7665-99-8**  
**(Cyclic GMP)**

L59 ANSWER 26 OF 30 MEDLINE on STN

ACCESSION NUMBER: 2006157240 MEDLINE Full-text

DOCUMENT NUMBER: PubMed ID: 16547056

TITLE: Control of formation and cellular detachment from  
Shewanella oneidensis MR-1 **biofilms** by cyclic  
di-GMP.AUTHOR: Thormann Kai M; Duttler Stefanie; Saville Renee M; Hyodo  
Mamoru; Shukla Soni; Hayakawa Yoshihiro; Spormann Alfred MCORPORATE SOURCE: Department of Civil Engineering, James H. Clark Center for  
Biomedical Engineering and Science, Stanford University,  
Stanford, CA 94305-5429, USA.SOURCE: Journal of bacteriology, (2006 Apr) Vol. 188, No. 7, pp.  
2681-91.

Journal code: 2985120R. ISSN: 0021-9193.

Report No.: NLM-PMC1428383.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
(RESEARCH SUPPORT, NON-U.S. GOV'T)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200604

ENTRY DATE: Entered STN: 21 Mar 2006

Last Updated on STN: 26 Apr 2006

Entered Medline: 25 Apr 2006

AB Stability and resilience against environmental perturbations are critical properties of medical and environmental **biofilms** and pose important targets for their control. **Biofilm** stability is determined by two mutually exclusive processes: attachment of cells to and detachment from the **biofilm** matrix. Using *Shewanella oneidensis* MR-1, an environmentally versatile, Fe(III) and Mn(IV) mineral-reducing microorganism, we identified mxdABCD as a new set of genes essential for formation of a three-dimensional **biofilm**. Molecular analysis revealed that mxdA encodes a cyclic bis(3',5')guanylic acid (cyclic di-GMP)-forming enzyme with an unusual GGDEF motif, i.e., NVDEF, which is essential for its function. mxdB encodes a putative **membrane** -associated glycosyl transferase. Both genes are essential for matrix attachment. The attachment-deficient phenotype of a  $\Delta$ mxdA mutant was rescued by ectopic expression of VCA0956, encoding another diguanylate cyclase. Interestingly, a rapid cellular detachment from the **biofilm** occurred upon induction of yhjH, a gene encoding an enzyme that has been shown to have phosphodiesterase activity. In this way, it was possible to bypass the previously identified sudden depletion of molecular oxygen as an environmental trigger to induce **biofilm** dissolution. We propose a model for **c-di-GMP** as a key intracellular regulator for **controlling biofilm** stability by shifting the state of a **biofilm** cell between attachment and detachment in a concentration-dependent manner.

TI Control of formation and cellular detachment from *Shewanella oneidensis* MR-1 **biofilms** by cyclic di-GMP.

AB Stability and resilience against environmental perturbations are critical properties of medical and environmental **biofilms** and pose important targets for their control. **Biofilm** stability is determined by two mutually exclusive

processes: attachment of cells to and detachment from the **biofilm** matrix. Using *Shewanella oneidensis* MR-1, an environmentally versatile, Fe(III) and Mn(IV) mineral-reducing microorganism, we identified mxdABCD as a new set of genes essential for formation of a three-dimensional **biofilm**. Molecular analysis revealed that mxdA encodes a cyclic bis(3',5')guanylic acid (cyclic di-GMP)-forming enzyme with an unusual GGDEF motif, i.e., NVDEF, which is essential for its function. mxdB encodes a putative **membrane** -associated glycosyl transferase. Both genes are essential for matrix attachment. The attachment-deficient phenotype of a DeltamxdA mutant was rescued by ectopic expression of VCA0956, encoding another diguanylate cyclase. Interestingly, a rapid cellular detachment from the **biofilm** occurred upon induction of yhjH, a gene encoding an enzyme that has been shown to have phosphodiesterase activity. In this way, it was possible to bypass the previously identified sudden depletion of molecular oxygen as an environmental trigger to induce **biofilm** dissolution. We propose a model for **c-di-GMP** as a key intracellular regulator for **controlling biofilm** stability by shifting the state of a **biofilm** cell between attachment and detachment in a concentration-dependent manner.

CT Bacterial Adhesion

\*Biofilms: GD, growth & development

\*Cyclic GMP: AA, analogs & derivatives

Cyclic GMP: ME, metabolism

Gene Expression Regulation, Bacterial

Operon

Polysaccharides: ME, metabolism

*Shewanella*: GE, genetics

\**Shewanella*: PH, physiology

*Shewanella*: UL, ultrastructure

RN 61093-23-0 (bis(3',5')-cyclic diguanylic acid); 7665-99-8  
(Cyclic GMP)

L59 ANSWER 27 OF 30 MEDLINE on STN

ACCESSION NUMBER: 2005389798 MEDLINE Full-text

DOCUMENT NUMBER: PubMed ID: 16048911

TITLE: 3',5'-cyclic

**diguanylic acid reduces the**  
**virulence of biofilm-forming**

***Staphylococcus aureus* strains in a mouse model of**  
**mastitis infection.**

AUTHOR: Brouillette Eric; Hyodo Mamoru; Hayakawa Yoshihiro;

Karaolis David K R; Malouin Francois

CORPORATE SOURCE: CEVDM, Departement de biologie, Faculte des sciences,  
Universite de Sherbrooke, 2500 Boul. Universite,  
Sherbrooke, Quebec, Canada.

SOURCE: Antimicrobial agents and chemotherapy, (2005 Aug) Vol. 49,  
No. 8, pp. 3109-13.  
Journal code: 0315061. ISSN: 0066-4804.  
Report No.: NLM-PMC1196217.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
(RESEARCH SUPPORT, NON-U.S. GOV'T)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200509

ENTRY DATE: Entered STN: 29 Jul 2005

Last Updated on STN: 20 Sep 2005

Entered Medline: 19 Sep 2005

AB The cyclic dinucleotide 3',5'-cyclic diguanylic acid (c-di-GMP) is a naturally occurring small molecule that regulates important signaling systems in bacteria. We have recently shown that **c-di -GMP inhibits *Staphylococcus aureus* biofilm** formation in vitro and its adherence to HeLa cells. We now

report that **c-di-GMP treatment** has an antimicrobial and antipathogenic activity in vivo and **reduces**, in a dose-dependent manner, **bacterial** colonization by **biofilm**-forming *S. aureus* strains in a mouse model of mastitis infection. Intramammary injections of 5 and 50 nmol of **c-di-GMP decreased colonization (bacterial CFU per gram of gland)** by 0.79 ( $P > 0.05$ ) and 1.44 ( $P < 0.01$ ) logs, respectively, whereas 200-nmol doses allowed clearance of the bacteria below the detection limit with a reduction of more than 4 logs ( $P < 0.001$ ) compared to the untreated control groups. These results indicate that cyclic dinucleotides potentially represent an attractive and novel drug platform which could be used alone or in combination with other agents or drugs in the **prevention, treatment, or control of infection**.

TI **3',5'-cyclic diguanylic**

**acid reduces** the virulence of **biofilm**-forming **Staphylococcus aureus** strains in a mouse model of mastitis infection.

AB The cyclic dinucleotide **3',5'-cyclic diguanylic acid (c-di-GMP)** is a naturally occurring small molecule that regulates important signaling systems in bacteria. We have recently shown that **c-di-GMP inhibits Staphylococcus aureus biofilm** formation in vitro and its adherence to HeLa cells. We now report that **c-di-GMP treatment** has an antimicrobial and antipathogenic activity in vivo and **reduces**, in a dose-dependent manner, **bacterial** colonization by **biofilm**-forming *S. aureus* strains in a mouse model of mastitis infection. Intramammary injections of 5 and 50 nmol of **c-di-GMP decreased colonization (bacterial CFU per gram of gland)** by 0.79 ( $P > 0.05$ ) and 1.44 ( $P < 0.01$ ) logs, respectively, whereas 200-nmol doses allowed clearance of the bacteria below the detection limit with a reduction of more than 4 logs ( $P < 0.001$ ) compared to the untreated control groups. These results indicate that cyclic dinucleotides potentially represent an attractive and novel drug platform which could be used alone or in combination with other agents or drugs in the **prevention, treatment, or control of infection**.

CT Check Tags: Female

Animals

Anti-Bacterial Agents: PD, pharmacology

\*Anti-Bacterial Agents: TU, therapeutic use

\*Biofilms: DE, drug effects

Biofilms: GD, growth & development

Cattle

\*Cyclic GMP: AA, analogs & derivatives

Cyclic GMP: PD, pharmacology

Cyclic GMP: TU, therapeutic use

\*Mastitis, Bovine: DT, drug therapy

Mastitis, Bovine: MI, microbiology

Mastitis, Bovine: PP, physiopathology

Mice

Models, Animal

Staphylococcal Infections: DT, drug therapy

Staphylococcal Infections: MI, microbiology

Staphylococcal Infections: PP, physiopathology

\*Staphylococcus aureus: DE, drug effects

Staphylococcus aureus: GD, growth & development

\*Staphylococcus aureus: PY, pathogenicity

Virulence

RN 61093-23-0 (bis(3',5')-cyclic diguanylic acid); 7665-99-8  
(Cyclic GMP)

L59 ANSWER 28 OF 30 MEDLINE on STN

ACCESSION NUMBER: 2002001765 MEDLINE [Full-text](#)

DOCUMENT NUMBER: PubMed ID: 11751251

TITLE: NO/cGMP signaling and HSP90 activity represses metamorphosis in the sea urchin *Lytechinus pictus*.

AUTHOR: Bishop C D; Brandhorst B P  
 CORPORATE SOURCE: Department of Molecular Biology and Biochemistry, Simon Fraser University, Burnaby, British Columbia V5A 1S6, Canada.  
 SOURCE: The Biological bulletin, (2001 Dec) Vol. 201, No. 3, pp. 394-404.  
 Journal code: 2984727R. ISSN: 0006-3185.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 (RESEARCH SUPPORT, NON-U.S. GOV'T)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200203  
 ENTRY DATE: Entered STN: 2 Jan 2002  
 Last Updated on STN: 3 Apr 2002  
 Entered Medline: 28 Mar 2002

- AB Nitric oxide (NO) signaling repressively regulates metamorphosis in two solitary ascidians and a gastropod. We present evidence for a similar role in the sea urchin *Lytechinus pictus*. NO commonly signals via soluble guanylyl cyclase (sGC). Nitric oxide synthase (NOS) activity in some **mammalian** cells, including neurons, depends on the molecular chaperone heat shock protein 90 (HSP90); this may be so in echinoid larvae as well. Pluteus larvae containing juvenile rudiments were **treated** with either radicicol L- or D-nitroarginine-methyl-ester (L-NAME and D-NAME), or IH-[1,2,4]oxadiazolo[4,3-a]quinoxalin-1-one (ODQ), inhibitors of HSP90, NOS, and sGC, respectively. In all instances, drug **treatment** significantly increased the frequency of metamorphosis. SNAP, a NO donor, suppressed the inductive properties of L-NAME and **biofilm**, a natural inducer of metamorphosis. NADPH diaphorase histochemistry indicated NOS activity in cells in the lower lip of the larval mouth, the preoral hood, the gut, and in the tube feet of the echinus rudiment. Histochemical staining coincided with NOS immunostaining. Microsurgical removal of the oral hood or the pre-oral hood did not induce metamorphosis, but larvae lacking these structures retained the capacity to metamorphose in response to ODQ. We propose that the production of NO repressively regulates the initiation of metamorphosis and that a sensory response to environmental cues reduces the production of NO, and consequently **cGMP**, to initiate metamorphosis.
- TI NO/**cGMP** signaling and HSP90 activity represses metamorphosis in the sea urchin *Lytechinus pictus*.
- AB Nitric oxide (NO) signaling repressively regulates metamorphosis in two solitary ascidians and a gastropod. We present evidence for a similar role in the sea urchin *Lytechinus pictus*. NO commonly signals via soluble guanylyl cyclase (sGC). Nitric oxide synthase (NOS) activity in some **mammalian** cells, including neurons, depends on the molecular chaperone heat shock protein 90 (HSP90); this may be so in echinoid larvae as well. Pluteus larvae containing juvenile rudiments were **treated** with either radicicol L- or D-nitroarginine-methyl-ester (L-NAME and D-NAME), or IH-[1,2,4]oxadiazolo[4,3-a]quinoxalin-1-one (ODQ), inhibitors of HSP90, NOS, and sGC, respectively. In all instances, drug **treatment** significantly increased the frequency of metamorphosis. SNAP, a NO donor, suppressed the inductive properties of L-NAME and **biofilm**, a natural inducer of metamorphosis. NADPH diaphorase histochemistry indicated NOS activity in cells in the lower lip of the larval mouth, the preoral hood, the gut, and in the tube feet of the echinus rudiment. Histochemical staining coincided with NOS immunostaining. Microsurgical removal of the oral hood or the pre-oral hood did not induce metamorphosis, but larvae lacking these structures retained the capacity to metamorphose in response to ODQ. We propose that the production of NO repressively regulates the initiation of metamorphosis and that a sensory response to environmental cues reduces the production of NO, and consequently **cGMP**, to initiate metamorphosis.
- CT Check Tags: Female; Male  
 Animals

**\*Cyclic GMP: PH, physiology**

- \*Enzyme Inhibitors: PD, pharmacology
    - Guanylate Cyclase: AI, antagonists & inhibitors
  - \*HSP90 Heat-Shock Proteins: PH, physiology
    - Lactones: PD, pharmacology
    - Macrolides
    - Metamorphosis, Biological: DE, drug effects
  - \*Metamorphosis, Biological: PH, physiology
    - NG-Nitroarginine Methyl Ester: PD, pharmacology
  - \*Nitric Oxide: PH, physiology
    - Nitric Oxide Donors: PD, pharmacology
    - Nitric Oxide Synthase: AI, antagonists & inhibitors
    - Oxadiazoles: PD, pharmacology
    - Quinoxalines: PD, pharmacology
    - S-Nitroso-N-Acetylpenicillamine: PD, pharmacology
    - Sea Urchins: DE, drug effects
  - \*Sea Urchins: GD, growth & development
    - Sea Urchins: PH, physiology
    - Signal Transduction: DE, drug effects
  - \*Signal Transduction: PH, physiology
- RN 10102-43-9 (Nitric Oxide); 12772-57-5 (monorden); 50903-99-6 (NG-Nitroarginine Methyl Ester); **7665-99-8 (Cyclic GMP)**; 79032-48-7 (S-Nitroso-N-Acetylpenicillamine)

L59 ANSWER 29 OF 30 EMBASE COPYRIGHT (c) 2009 Elsevier B.V. All rights reserved on STN

ACCESSION NUMBER: 2009247317 EMBASE Full-text  
 TITLE: In vivo evaluation of vaginal films for **mucosal** delivery of nitric oxide.  
 AUTHOR: Yoo, Jin-Wook; Acharya, Gayathri; Lee, Chi H. (correspondence)  
 CORPORATE SOURCE: Division of Pharmaceutical Sciences, School of Pharmacy, University of Missouri at Kansas City, 5005 Rockhill Rd, MO 64110, United States. leech@umkc.edu  
 SOURCE: Biomaterials, (August 2009) Vol. 30, No. 23-24, pp. 3978-3985.  
 Refs: 38  
 ISSN: 0142-9612 CODEN: BIMADU  
 PUBLISHER: Elsevier Ltd, Langford Lane, Kidlington, Oxford, OX5 1GB, United Kingdom.  
 PUBLISHER IDENT.: S 0142-9612(09)00350-0  
 COUNTRY: United Kingdom  
 DOCUMENT TYPE: Journal; Article  
 FILE SEGMENT: 037 Drug Literature Index  
 039 Pharmacy  
 LANGUAGE: English  
 SUMMARY LANGUAGE: English  
 ENTRY DATE: Entered STN: 16 Jun 2009  
 Last Updated on STN: 16 Jun 2009

AB Nitric oxide (NO)-releasing vaginal films were developed and evaluated as a potential advanced treatment option for female sexual arousal disorder (FSAD). The polymeric films containing s-nitrosoglutathione (GSNO), an endogenous NO donor, were prepared using the reduced-pressure drying method. The surface morphology, thermal/mechanical properties, stability, loading efficiency and physico-dynamic properties were characterized and the pharmacological activities were evaluated through in vitro and in vivo studies. The GSNO films were homogeneous and transparent, and showed suitable mucoadhesiveness and mechanical properties. The release profiles of NO from the GSNO films followed the first-order kinetic pattern and NO activated the NO-**cGMP** signaling pathway in vaginal cells. The GSNO films significantly enhanced the



duration of action of GSNO and vagina blood perfusion in the rat model without causing any cytotoxic effects. The NO-releasing vaginal films might be used as a promising treatment device against FSAD.

- TI In vivo evaluation of vaginal films for **mucosal** delivery of nitric oxide.
- AB Nitric oxide (NO)-releasing vaginal films were developed and evaluated as a potential advanced treatment option for female sexual arousal disorder (FSAD). The polymeric films containing s-nitrosoglutathione (GSNO), an endogenous NO donor, were prepared using the reduced-pressure drying method. The surface morphology, thermal/mechanical properties, stability, loading efficiency and physico-dynamic properties were characterized and the pharmacological activities were evaluated through in vitro and in vivo studies. The GSNO films were homogeneous and transparent, and showed suitable mucoadhesiveness and mechanical properties. The release profiles of NO from the GSNO films followed the first-order kinetic pattern and NO activated the NO-**cGMP** signaling pathway in vaginal cells. The GSNO films significantly enhanced the duration of action of GSNO and vagina blood perfusion in the rat model without causing any cytotoxic effects. The NO-releasing vaginal films might be used as a promising treatment device against FSAD.
- CT Medical Descriptors:  
 animal experiment  
 article  
     **\*biofilm**  
 cell type  
 controlled study  
 cytotoxicity  
 \*drug delivery system  
 drug release  
 drug stability  
 female  
 female sexual dysfunction  
 human  
 human cell  
 in vitro study  
 in vivo study  
 morphology  
 mucoadhesion  
 nonhuman  
 perfusion  
 priority journal  
 rat  
 sexual arousal disorder  
 vagina cell  
 \*vaginal film
- CT Drug Descriptors:  
 carbomer: PR, pharmaceuticals  
     **cyclic GMP: EC, endogenous compound**  
 hydroxypropylmethylcellulose: PR, pharmaceuticals  
 macrogol: PR, pharmaceuticals  
 methylcellulose  
 \*nitric oxide: PR, pharmaceuticals  
 s nitrosoglutathione: PR, pharmaceuticals  
 s nitrosoglutathione: TP, topical drug administration
- RN (carbomer) 9007-20-9, 9062-04-8; (**cyclic GMP**)  
**7665-99-8**; (hydroxypropylmethylcellulose) 9004-65-3; (macrogol)  
 25322-68-3; (methylcellulose) 79484-92-7, 9004-67-5; (nitric oxide)  
 10102-43-9; (s nitrosoglutathione) 57564-91-7

ACCESSION NUMBER: 2009280434 EMBASE Full-text  
 TITLE: Signals, regulatory networks, and materials that build and break bacterial **biofilms**.  
 AUTHOR: Karatan, Ece (correspondence)  
 CORPORATE SOURCE: Department of Biology, Appalachian State University, Boone, NC 28608, United States. karatane@appstate.edu  
 AUTHOR: Watnick, Paula  
 CORPORATE SOURCE: Division of Infectious Diseases, Children's Hospital, Boston, MA 02115, United States.  
 AUTHOR: Watnick, Paula  
 CORPORATE SOURCE: Department of Microbiology and Molecular Genetics, Harvard Medical School, Boston, MA 02115, United States.  
 SOURCE: Microbiology and Molecular Biology Reviews, (June 2009) Vol. 73, No. 2, pp. 310-347.  
 Refs: 363  
 ISSN: 1092-2172 CODEN: MMBRF7  
 PUBLISHER: American Society for Microbiology, 1752 N Street N.W., Washington, DC 20036-2904, United States.  
 COUNTRY: United States  
 DOCUMENT TYPE: Journal; General Review; (Review)  
 FILE SEGMENT: 004 Microbiology: Bacteriology, Mycology, Parasitology and Virology  
 LANGUAGE: English  
 ENTRY DATE: Entered STN: 7 Jul 2009  
 Last Updated on STN: 7 Jul 2009  
 TI Signals, regulatory networks, and materials that build and break bacterial **biofilms**.  
 CT Medical Descriptors:  
 antibiotic resistance  
 bacterial flagellum  
**\*bacterial membrane**  
 bacterium pilus  
**\*biofilm**  
 Caulobacter crescentus  
 materials testing  
 molecular biology  
 nonhuman  
 Pseudomonas aeruginosa  
 Pseudomonas fluorescens  
 regulatory mechanism  
 review  
 Salmonella typhimurium  
 second messenger  
 signal transduction  
 transcription regulation  
**Vibrio cholerae**  
**Vibrio harveyi**  
 Yersinia enterocolitica  
 Yersinia pseudotuberculosis  
 CT Drug Descriptors:  
 \*adhesin: EC, endogenous compound  
 antiinfective agent  
 cyclic AMP: EC, endogenous compound  
**cyclic GMP: EC, endogenous compound**  
 exopolysaccharide  
 glucose  
 iron  
 monosaccharide  
 tobramycin  
 RN (cyclic AMP) 60-92-4; (cyclic GMP) 7665-99-8